

1 **ON THE EXTINCTION OF MULTIPLE SHOCKS**
2 **IN SCALAR VISCIOUS CONSERVATION LAWS**

3 JEANNE LIN, DMITRY E. PELINOVSKY, AND BJÖRN DE RIJK

ABSTRACT. We are interested in the dynamics of interfaces, or zeros, of shock waves in general scalar viscous conservation laws with a locally Lipschitz continuous flux function, such as the modular Burgers' equation. We prove that all interfaces coalesce within finite time, leaving behind either a single interface or no interface at all. Our proof relies on mass and energy estimates, regularization of the flux function, and an application of the Sturm theorems on the number of zeros of solutions of parabolic problems. Our analysis yields an explicit upper bound on the time of extinction in terms of the initial condition and the flux function. Moreover, in the case of a smooth flux function, we characterize the generic bifurcations arising at a coalescence event with and without the presence of odd symmetry. We identify associated scaling laws describing the local interface dynamics near collision. Finally, we present an extension of these results to the case of anti-shock waves converging to asymptotic limits of opposite signs. Our analysis is corroborated by numerical simulations in the modular Burgers' equation and its regularizations.

4 1. INTRODUCTION

We consider shock and anti-shock waves with multiple interfaces in the scalar viscous conservation law

$$u_t = u_{xx} + f(u)_x, \quad t \geq 0, \quad x \in \mathbb{R}, \quad u(t, x) \in \mathbb{R}, \quad (1.1)$$

5 where $f: \mathbb{R} \rightarrow \mathbb{R}$ is a locally Lipschitz continuous flux function. A classical example is
6 the viscous Burgers' equation with $f(u) = u^2$. Our regularity assumption on f allows for
7 nonsmooth choices such as $f(u) = |u|$, yielding the modular Burgers' equation which has
8 been used to model inelastic dynamics of particles with piecewise interaction potentials [7,
9 16] and whose behavior has been studied analytically and numerically in [9, 13, 15].

Shock waves are solutions of (1.1) with initial data $u_0(x)$ converging to nonzero asymptotic limits ϕ_{\pm} as $x \rightarrow \pm\infty$, which satisfy $\phi_+ \neq \phi_-$ and obey the Gel'fand-Oleinik entropy condition

$$\frac{f(\phi_{\min}) - f(z)}{z - \phi_{\min}} > \frac{f(\phi_-) - f(\phi_+)}{\phi_+ - \phi_-}, \quad z \in (\phi_{\min}, \phi_{\max}), \quad (1.2)$$

10 where we denote $\phi_{\min} = \min\{\phi_-, \phi_+\}$ and $\phi_{\max} = \max\{\phi_-, \phi_+\}$. On the other hand, *anti-*
11 *shock waves* are solutions of (1.1) with initial data $u_0(x)$ converging to nonzero asymptotic
12 limits ϕ_{\pm} , which satisfy $\phi_+ \neq \phi_-$ and do not fulfill the entropy condition (1.2).

The Gel'fand-Oleinik entropy condition (1.2) is consistent with the existence of *traveling shock waves*, which are solutions of (1.1) of the form $u(t, x) = \phi(x - ct)$, where $c \in \mathbb{R}$ denotes the propagation speed and the profile $\phi: \mathbb{R} \rightarrow \mathbb{R}$ solves the scalar problem

$$0 = \phi_\xi + c(\phi - \phi_-) + f(\phi) - f(\phi_-).$$

Here, the profile $\phi(\xi)$ converges to the asymptotic limits ϕ_\pm as $\xi \rightarrow \pm\infty$ and the speed is given by the Rankine-Hugoniot condition

$$c = \frac{f(\phi_-) - f(\phi_+)}{\phi_+ - \phi_-}. \quad (1.3)$$

13 The traveling shock-wave solution $u(t, x) = \phi(x - ct)$ defined for $\phi_- \neq \phi_+$ exists if and
14 only if the entropy condition (1.2) is fulfilled.

15 Traveling shock waves form an important class of asymptotic solutions of (1.1) in
16 the sense that they serve as global attractors for shock waves. More precisely, for twice
17 continuously differentiable flux functions f , it has been proven in [4, 8] that any shock-
18 wave solution of the viscous conservation law (1.1) converges as $t \rightarrow \infty$ in both L^1 - and
19 L^∞ -norm to a traveling shock wave, which necessarily possesses the same asymptotic
20 limits ϕ_\pm at $\pm\infty$.

21 We are interested in the temporal dynamics of zeros, so-called *interfaces*, for shock-
22 and anti-shock wave solutions of the viscous conservation law (1.1). In our analysis we
23 distinguish between three classes of initial data u_0 , where both ϕ_+ and ϕ_- are nonzero:

- 24 • **Class I:** $u_0(x)$ converges to asymptotic limits ϕ_\pm of *opposite signs* as $x \rightarrow \pm\infty$,
25 which obey the Gel'fand-Oleinik entropy condition (1.2);
- 26 • **Class II:** $u_0(x)$ converges to asymptotic limits ϕ_\pm of the *same sign* as $x \rightarrow \pm\infty$;
- 27 • **Class III:** $u_0(x)$ converges to asymptotic limits ϕ_\pm of *opposite signs* as $x \rightarrow \pm\infty$,
28 which do *not* satisfy the entropy condition (1.2).

29 We note that solutions of (1.1) with initial data of the class I are shock waves, whereas
30 solutions of (1.1) with initial data of class III are anti-shock waves. Although solutions
31 of (1.1) with initial data of class II can be either shock or anti-shock waves, it is not
32 necessary to distinguish between them in our analysis.

33 In addition to the above assumptions, we require that our initial datum u_0 is uniformly
34 continuous and bounded, and that $u_0 - \phi_\pm$ is L^1 -integrable on \mathbb{R}_\pm . Then, by the com-
35 parison principle and standard parabolic regularity theory [10], the solution of (1.1) with
36 initial condition u_0 stays bounded and is continuously differentiable for all positive times,
37 while maintaining its asymptotic limits ϕ_\pm at $\pm\infty$. Nevertheless, if the flux function f
38 is not continuously differentiable, as in the case of the modular Burgers' equation, the
39 second derivative of the solution of (1.1) may be discontinuous [9].

40 The classical Sturm Theorems yield that in parabolic semilinear equations the number
41 of zeros of solutions is nonincreasing over time. Moreover, if at some time t_0 the solution
42 has an (isolated) multiple zero x_0 , then, in a sufficiently small neighborhood of x_0 , the

43 number of zeros strictly decreases when t passes through t_0 . We refer to [5] for a survey
44 on Sturm's Theorems and their applications.

45 In this paper we study *finite-time coalescence* of interfaces. In preliminary work [13]
46 we showed that the evolution of odd shock waves with three symmetric interfaces in
47 the modular Burgers' equation leads to a finite-time coalescence of these interfaces to a
48 single interface and we conjectured a scaling law for the local interface dynamics near the
49 collision event based on data fitting. In this work we extend these results to general viscous
50 conservation laws of the form (1.1) and establish finite-time coalescence of interfaces for all
51 solutions with initial data of class I or II and thus, of all shock-wave solutions. Moreover,
52 we show that in the specific case of the modular Burgers' equation, solutions with initial
53 data of class III, i.e. anti-shock waves, can also exhibit finite-time coalescence of interfaces.

54 For solutions of (1.1) with initial data of class I, we establish that all interfaces must co-
55 alesce to a single interface within finite time. The argument generalizes the idea from [13]
56 and relies on a differential inequalities for the masses of $u(t, x) - \phi_+$ and $u(t, x) - \phi_-$
57 measured with respect to the position of the interface, in combination with smooth ap-
58 proximation of the flux function and an application of the Sturm Theorem from [1]. Our
59 analysis yields an explicit upper bound on the time at which all interfaces have collapsed
60 to a single interface. We emphasize that although the results in [4, 8] imply that solutions
61 with initial data of class I converge in L^1 - and L^∞ -norms to a traveling shock wave, which
62 must necessarily be strictly monotone and thus, has precisely a single interface, this is not
63 sufficient to conclude finite-time coalescence to a single interface because interfaces of the
64 solution might accumulate close to the interface of the associated traveling shock wave.

65 Initial data u_0 of class II can always be bounded from above or below by a smooth
66 function \tilde{u}_0 , which satisfies $\tilde{u}_0(x) \rightarrow \tilde{u}_\infty$ as $x \rightarrow \pm\infty$, where $\tilde{u}_\infty \neq 0$ has the same sign
67 as ϕ_\pm . For twice continuously differentiable flux functions f , the finite-time extinction of
68 all interfaces of the solution $\tilde{u}(t, \cdot)$ of (1.1) with initial condition \tilde{u}_0 follows by evoking
69 the result from [4] that $\tilde{u}(t, \cdot)$ converges in L^∞ -norm to the constant state \tilde{u}_∞ as $t \rightarrow \infty$.
70 Consequently, the comparison principle yields the finite-time extinction of interfaces of
71 the solution $u(t, \cdot)$ of (1.1) with initial condition u_0 . Yet, the result in [4] does not provide
72 an explicit upper bound on the extinction time and does not readily apply to the current
73 setting of locally Lipschitz continuous flux functions. To extend the conclusion to our
74 setting, we apply a softer argument based on energy estimates, smooth approximation of
75 the flux function, conservation of mass and the Gagliardo-Nirenberg inequality to yield
76 an explicit upper bound on the time at which all interfaces of $\tilde{u}(t, \cdot)$, and thus, also of
77 $u(t, \cdot)$, have gone extinct, cf. Remark 3.6.

78 Whether solutions of (1.1) with initial data of class III do exhibit finite-time coales-
79 cence of interfaces to a single interface is currently an open problem. Since the entropy
80 condition (1.2) is not fulfilled, there exists no traveling shock to which $u(t, \cdot)$ can converge
81 in norm as $t \rightarrow \infty$. To shed some light on this open question, we consider anti-shock
82 waves with initial data of class III in the modular Burgers' equation with flux function
83 $f(u) = |u|$. Our analysis indicates that, although all interfaces coalesce to a single inter-
84 face in this case, the anti-shock wave converges *locally uniformly* to 0 as $t \rightarrow \infty$ suggesting

85 that obtaining a result in general might be subtle or even false. Colloquially speaking,
 86 since the solution profile can converge to 0 uniformly, locally near interfaces, diffusion
 87 might be too weak to enforce coalescence of interfaces. In fact, recent results [6] imply
 88 that the ω -limit set (in the locally uniform topology induced by $L_{\text{loc}}^\infty(\mathbb{R})$) of bounded
 89 solutions of scalar viscous conservation laws (1.1) can be complicated in the sense that
 90 it can contain a solution that is neither a traveling shock nor a constant, underlining a
 91 fundamental difference between shock waves and general bounded solutions of (1.1).

92 In addition to establishing finite-time coalescence of interfaces of shock and anti-shock
 93 waves, we study the interface dynamics about a coalescence event in the case of a smooth
 94 flux function f . If a coalescence event occurs for a solution $u(t, x)$ of (1.1) at some time
 95 $t = t_0$ and point $x = \xi_0$, it must hold that $u_x(t_0, \xi_0) = 0$ and it follows from one of the
 96 classical Sturm Theorems [1] that there exist $\delta > 0$ and a neighborhood $U \subset \mathbb{R}$ of ξ_0
 97 such that for $t \in (t_0 - \delta, t_0)$, there are at least two interfaces in U and for $t \in (t_0, t_0 + \delta)$,
 98 there is at most one interface in U . Without the presence of additional symmetries, one
 99 generically has $u_{xx}(t_0, \xi_0) \neq 0$. We show that in this situation a *fold bifurcation* occurs.
 100 That is, there are precisely two interfaces $\xi_1(t) < \xi_2(t)$ in U for $t \in (t_0 - \delta, t_0)$ and no
 101 interfaces in U for $t \in (t_0, t_0 + \delta)$. Moreover, we obtain the scaling law

$$102 \quad \xi_{1,2}(t) - \xi_0 \sim \pm \sqrt{2(t_0 - t)} \quad \text{as } t \rightarrow t_0^-. \quad (1.4)$$

103 In the case of an odd reflection symmetry, we generically have $u_{xx}(t_0, \xi_0) = 0$ and
 104 $u_{xxx}(t_0, \xi_0) \neq 0$. This leads to a *pitchfork bifurcation*, for which there are precisely three
 105 interfaces $\xi_1(t) < \xi(t) < \xi_2(t)$ in U for $t \in (t_0 - \delta, t_0)$ and exactly one interface $\xi(t)$ remains
 106 in U for $t \in (t_0, t_0 + \delta)$. We also identify the associated scaling laws

$$107 \quad \xi_{1,2}(t) - \xi_0 \sim \pm \sqrt{6(t_0 - t)} \quad \text{as } t \rightarrow t_0^- \quad (1.5)$$

108 and

$$109 \quad \xi(t) - \xi_0 \sim \alpha(t_0 - t) \quad \text{as } t \rightarrow t_0 \quad (1.6)$$

110 for some $\alpha \in \mathbb{R}$. We show that the conditions for a pitchfork bifurcation are satisfied in
 111 the classical Burgers' equation with flux function $f(u) = u^2$ for odd shock waves with
 112 a single zero on $(0, \infty)$. We note that the above results yield that the lower and upper
 113 bounds in the Sturm Theorem [1, Theorem B] on the number of interfaces before and
 114 after a coalescence event are sharp.

115 Finally, we corroborate our results with numerical simulations of the modular Burgers'
 116 equation. Our numerical approximations rely on a regularization of the modular
 117 nonlinearity and employ an elementary finite-difference scheme. These numerical approx-
 118 imations are different from those used in [13], where the modular Burgers' equation was
 119 solved on a partition of a real line complemented with additional boundary conditions at
 120 the interfaces. We study odd shock and anti-shock waves and observe finite-time coales-
 121 cence of interfaces through a pitchfork bifurcation. In addition, the numerics confirms
 122 the same scaling law (1.5) for the interface extinction.

123 The derivation of scaling laws describing the interface dynamics near coalescence has
 124 been addressed in other contexts as well and appeared to be challenging. In [2] a linear
 125 inhomogeneous heat equation was considered as a simple model for oxygen diffusion. It

126 was suggested that the oxygen front (the interface) collapses according to the scaling law
 127 $(t_0 - t)^{1/2}$. However, a more recent study in [12] based on new numerical algorithms for the
 128 time-dependent Stefan problem showed that the scaling law $(t_0 - t)^{1/2}$ is not accurate due
 129 to an additional singularity as $t \rightarrow t_0^-$. Other interface models were studied in [18, 19]
 130 by means of matched asymptotic expansions in the context of a KPP equation with a
 131 discontinuous cut-off in the reaction function.

132 We conjecture that the scaling laws (1.4), (1.5), and (1.6) proven for smooth flux
 133 functions remain true for locally Lipschitz continuous flux functions such as the modular
 134 Burgers' equation. However, this question remains open for future research.

135 This paper is organized as follows. In Section 2 we state well-posedness and approxi-
 136 mation results for solutions of the viscous conservation law (1.1). Section 3 is devoted to
 137 the analysis of finite-time coalescence of interfaces for solutions with initial data of class
 138 I, II, and III. In Section 4 we analyze the fold and pitchfork bifurcations describing the
 139 interface dynamics near coalescence events and derive associated scaling laws. Section 5
 140 presents numerical simulations illustrating the pitchfork bifurcation for both shock and
 141 anti-shock waves in a regularized version of the modular Burgers' equation. Appendix A
 142 contains the proofs of the well-posedness and approximation results of Section 2.

143 **Acknowledgements.** J. Lin was supported by Stewart Research Scholarship of Mc-
 144 Master University. D. E. Pelinovsky acknowledges the funding of this study provided by
 145 the grant No. FSWE-2023-0004 and grant No. NSH-70.2022.1.5.

146 2. GLOBAL WELL-POSEDNESS AND APPROXIMATION

In this section we establish global well-posedness of uniformly continuous and bounded
 solutions of the viscous conservation law (1.1). We first consider smooth flux functions
 f before studying the general case of a locally Lipschitz continuous flux function. We
 show that by locally approximating the flux function f by a smooth function \tilde{f} , one can
 approximate solutions $u(t, \cdot)$ of (1.1) on any finite time interval by a solution $\tilde{u}(t, \cdot)$ of the
 regularized problem

$$\tilde{u}_t = \tilde{u}_{xx} + \tilde{f}(\tilde{u})_x. \quad (2.1)$$

147 Proofs of all results formulated in this section can be found in Appendix A.

148 For smooth flux functions $f \in C^\infty(\mathbb{R})$ local existence and uniqueness of classical
 149 solutions of (1.1) follow readily by standard regularity theory for parabolic semilinear
 150 equations [10]. The fact that (1.1) obeys a comparison principle [14, 17] then yields
 151 global well-posedness. All in all, we establish the following result.

Lemma 2.1. *Let $f \in C^\infty(\mathbb{R})$ and $u_0 \in C_{\text{ub}}^1(\mathbb{R})$. Let $M_0 = \sup\{u_0(x) : x \in \mathbb{R}\}$ and
 $m_0 = \inf\{u_0(x) : x \in \mathbb{R}\}$. There exists a unique smooth global classical solution*

$$u \in C([0, \infty), C_{\text{ub}}^1(\mathbb{R})) \cap C((0, \infty), C_{\text{ub}}^2(\mathbb{R})) \cap C^1((0, \infty), C_{\text{ub}}(\mathbb{R})),$$

152 of (1.1) with initial condition $u(0, \cdot) = u_0$ such that $m_0 \leq u(t, x) \leq M_0$ for all $t \geq 0$ and
 153 $x \in \mathbb{R}$. Moreover, we have $u \in C^\infty((0, \infty) \times \mathbb{R}, \mathbb{R})$ with $\partial_t^k u(t, \cdot) \in C_{\text{ub}}^l(\mathbb{R})$ for $t \geq 0$ and
 154 $k, l \in \mathbb{N}_0$.

Next, we establish global well-posedness of solutions of (1.1) for locally Lipschitz continuous flux functions f . In this case, classical solutions in the sense of Lemma 2.1 cannot always be expected. For instance, the modular Burgers' equation with flux function $f(u) = |u|$ admits for any $\phi_\pm \in \mathbb{R}$ with $\phi_- < 0 < \phi_+$ a traveling shock-wave solution $u(t, x) = \phi(x - ct)$ converging to asymptotic limits ϕ_\pm and propagating with speed

$$c = \frac{\phi_+ + \phi_-}{\phi_- - \phi_+},$$

whose profile

$$\phi(\pm\xi) = \phi_\pm (1 - e^{-(1+c)\xi}), \quad \xi \geq 0,$$

does lie in $C_{\text{ub}}^1(\mathbb{R})$, but not in $C_{\text{ub}}^2(\mathbb{R})$. Therefore, we consider *mild* solutions of (1.1), which solve the associated integral equation

$$u(t, \cdot) = e^{\partial_x^2 t} u_0 + \int_0^t \partial_x e^{\partial_x^2(t-s)} f(u(s, \cdot)) ds, \quad (2.2)$$

155 where $u(0, \cdot) = u_0 \in C_{\text{ub}}^1(\mathbb{R})$ denotes the initial condition.

156 Standard analytic semigroup theory in combination with the fact that f is locally
 157 Lipschitz continuous yields local existence and uniqueness of solutions of (2.2) in $C_{\text{ub}}(\mathbb{R})$.
 158 We note that it is important here to compose the derivative in (2.2) with the semigroup
 159 $e^{\partial_x^2(t-s)}$, rather than applying it to the flux function f , since f' is not necessarily locally
 160 Lipschitz continuous. Global well-posedness follows by approximating the solution $u(t, \cdot)$
 161 of (2.2) by the global classical solution $\tilde{u}(t, \cdot)$ of the regularized problem (2.1), where
 162 $\tilde{f} \in C^\infty(\mathbb{R})$ is a smooth local approximation of f . This leads to the following result.

Lemma 2.2. *Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be locally Lipschitz continuous and $u_0 \in C_{\text{ub}}^1(\mathbb{R})$. Let $M_0 = \sup\{u_0(x) : x \in \mathbb{R}\}$ and $m_0 = \inf\{u_0(x) : x \in \mathbb{R}\}$. There exists a unique global solution $u \in C([0, \infty), C_{\text{ub}}(\mathbb{R}))$ of (2.2) such that $m_0 \leq u(t, x) \leq M_0$ for all $t \geq 0$ and $x \in \mathbb{R}$. Moreover, there exist constants $\delta_0, C_0 > 0$ such that for each $\delta \in (0, \delta_0)$ and $\tilde{f} \in C^\infty(\mathbb{R})$ satisfying*

$$\sup \left\{ |f(v) - \tilde{f}(v)| : v \in [m_0, M_0] \right\} < \delta,$$

the global classical solution

$$\tilde{u} \in C([0, \infty), C_{\text{ub}}^1(\mathbb{R})) \cap C((0, \infty), C_{\text{ub}}^2(\mathbb{R})) \cap C^1((0, \infty), C_{\text{ub}}(\mathbb{R})), \quad (2.3)$$

of the regularized equation (2.1) with $\tilde{u}(0, \cdot) = u_0$, established in Lemma 2.1, obeys the estimates

$$m_0 \leq \tilde{u}(t, x) \leq M_0, \quad \|u(t, \cdot) - \tilde{u}(t, \cdot)\|_\infty \leq C_0 \delta \sqrt{t}, \quad (2.4)$$

163 for all $t \geq 0$ and $x \in \mathbb{R}$.

164 Next, we approximate mild solutions of (1.1) by solutions of the regularized equa-
 165 tion (2.1) in C_{ub}^1 -norm rather than in C_{ub} -norm. The approximation in C_{ub}^1 -norm will
 166 be used in the upcoming analysis to conclude that a single interface of the approximate
 167 solution also yields a single interface of the original solution.

Lemma 2.3. *Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be locally Lipschitz continuous. Let $u \in C([0, \infty), C_{\text{ub}}^1(\mathbb{R}))$ be a global solution of (2.2) with initial condition $u(0) = u_0 \in C_{\text{ub}}^1(\mathbb{R})$. Set $M_0 = \sup\{u_0(x) : x \in \mathbb{R}\}$ and $m_0 = \inf\{u_0(x) : x \in \mathbb{R}\}$. Let $R, \tau, \varepsilon > 0$. There exists $\delta_0 > 0$ such that for each $\delta \in (0, \delta_0)$ and $\tilde{f} \in C^\infty(\mathbb{R})$ satisfying*

$$\sup \left\{ |f(v) - \tilde{f}(v)| : v \in [m_0, M_0] \right\} < \delta, \quad \sup \left\{ |\tilde{f}'(v)| : v \in [m_0, M_0] \right\} \leq R,$$

the global classical solution (2.3) of the regularized equation (2.1) with initial condition $\tilde{u}(0, \cdot) = u_0$, established in Lemma 2.1, obeys the estimates

$$m_0 \leq \tilde{u}(t, x) \leq M_0 \quad \sup_{0 \leq s \leq \tau} \|u(s, \cdot) - \tilde{u}(s, \cdot)\|_{W^{1, \infty}} < \varepsilon, \quad (2.5)$$

168 for $x \in \mathbb{R}$ and $t \geq 0$.

169 We emphasize that Lemma 2.3, in contrast to Lemma 2.2, is merely an approximation
 170 result and does not imply the existence of a global mild solution in $C_{\text{ub}}^1(\mathbb{R})$. This suffices
 171 for our purposes because we only apply Lemma 2.3 to establish finite-time coalescence
 172 of interfaces for solutions of (1.1) with initial data of class I, for which global existence
 173 of a mild solution in $C_{\text{ub}}^1(\mathbb{R})$ follows from a separate well-posedness result, which we will
 174 formulate next.

175 In case of initial data of class I the entropy condition (1.2) yields the existence of
 176 a traveling shock wave with the same limits at $\pm\infty$. We require that the difference
 177 between the initial condition and the traveling shock wave is L^1 -integrable and show
 178 that this integrability is maintained over time, which will be important for the mass and
 179 energy estimates in the upcoming proofs establishing finite-time coalescence of interfaces
 180 in §3. Moreover, by integrating the viscous conservation law (1.1) we obtain global well-
 181 posedness of mild solutions in $C_{\text{ub}}^1(\mathbb{R})$ rather than in $C_{\text{ub}}(\mathbb{R})$.

Lemma 2.4. *Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be locally Lipschitz continuous and let $u_0 \in C_{\text{ub}}^1(\mathbb{R})$. Suppose that there exist $c, C \in \mathbb{R}$ and a solution $\phi \in C_{\text{ub}}^1(\mathbb{R})$ of the profile equation*

$$0 = \phi_\xi + c\phi + f(\phi) + C.$$

182 *Suppose $u_0 - \phi$ is L^1 -integrable. Then, there exists a unique solution $u \in C([0, \infty), C_{\text{ub}}^1(\mathbb{R}))$
 183 of (2.2) such that $u(t, \cdot) - \phi$ is L^1 -integrable for all $t \geq 0$.*

184 3. FINITE-TIME COALESCENCE OF INTERFACES

185 Here we establish finite-time coalescence of interfaces for solutions $u(t, \cdot)$ of (1.1) with
 186 initial data $u(0, \cdot) = u_0 \in C_{\text{ub}}^1(\mathbb{R})$ of class I or II. We emphasize that solutions with such
 187 initial data include all shock waves. On the other hand, anti-shock waves converging to
 188 asymptotic limits of opposite signs are not included. We study finite-time coalescence of

189 interfaces of this type of anti-shock waves at the end of this section in the specific setting
190 of the modular Burgers' equation.

191 **3.1. Solutions with initial data of class I.** Observing that solutions $u(t, x)$ of (1.1)
192 with initial data $u(0, \cdot) = u_0 \in C_{\text{ub}}^1(\mathbb{R})$ of type I maintain their asymptotic limits ϕ_{\pm} as
193 $x \rightarrow \pm\infty$ for every $t > 0$ by Lemma 2.4, it readily follows that the solution possesses at
194 least one interface for all $t \geq 0$ since ϕ_+ and ϕ_- have opposite signs. We establish that
195 all interfaces coalesce to a single one within finite time in this case.

196 **Theorem 3.1.** *Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be locally Lipschitz continuous and $u_0 \in C_{\text{ub}}^1(\mathbb{R})$. Suppose*
197 *$u_0(x)$ converges to asymptotic limits ϕ_{\pm} as $x \rightarrow \pm\infty$ such that ϕ_+ and ϕ_- have opposite*
198 *signs and the Gel'fand-Oleinik entropy condition (1.2) holds. Moreover, assume that*
199 *$u_0 - \phi_{\pm}$ is L^1 -integrable on \mathbb{R}_{\pm} and we have $u_0(x) \in [\min\{\phi_-, \phi_+\}, \max\{\phi_-, \phi_+\}]$ for all*
200 *$x \in \mathbb{R}$. Let $u \in C([0, \infty), C_{\text{ub}}^1(\mathbb{R}))$ be the global mild solution of (1.1), established in*
201 *Lemma 2.4. Then, there exists a time $T > 0$ such that for all $t > T$ the solution $u(t, \cdot)$*
202 *possesses precisely one zero.*

203 The proof of Theorem 3.1 is based on ideas developed in [13], where it is shown that
204 the interfaces of odd shock waves in the modular Burgers' equation coalesce to a single
205 one within finite time. The analysis in [13] relies on a differential inequality for the mass
206 measured with respect to the fixed interface at 0. Indeed, due to odd symmetry, 0 is
207 necessarily an interface of the shock wave for all time and must be the middle interface.

In the general setting considered here, without the presence of an odd symmetry,
interfaces are a priori not fixed, which suggests mass functions of the form

$$\mathcal{M}_1(t) = \int_{-\infty}^{\xi_2(t)} (u(t, x) - \phi_-) dx, \quad \mathcal{M}_2(t) = \int_{\xi_2(t)}^{\infty} (\phi_+ - u(t, x)) dx, \quad (3.1)$$

208 where $\xi_2(t)$ is an interface of $u(t, \cdot)$, which now depends on time. As in [13] we aim to show
209 that the assumption that $\xi_2(t)$ is an interface lying strictly in between two other interfaces
210 $\xi_1(t), \xi_3(t)$ leads to a contradiction with certain inequalities obeyed by the mass functions
211 $\mathcal{M}_1(t)$ and $\mathcal{M}_2(t)$. This then yields an explicit time $T > 0$ such that $\xi_1(t) < \xi_2(t) < \xi_3(t)$
212 cannot hold for $t > T$.

213 To derive the desired inequalities for $\mathcal{M}_1(t)$ and $\mathcal{M}_2(t)$, a standard strategy is to differ-
214 entiate with respect to time (using the Leibniz' integral rule) and use the equation (1.1) to
215 express temporal derivatives of $u(t, x)$. Yet, as mentioned in §2, it cannot be expected in
216 the case of a locally Lipschitz continuous flux function f that $u(t, x)$ is a classical solution
217 of (1.1), which is differentiable with respect to time and twice differentiable with respect
218 to space. In addition, even if the flux function f were smooth, the interface $\xi_2(t)$, being
219 a root of the C^1 -function $u(t, x)$, is not necessarily differentiable. In fact, the upcoming
220 analysis in §4 shows that $\xi_2(t)$ may fail to be differentiable if two interfaces collide.

221 To address the first challenge we approximate the solution $u(t, x)$ of (1.1) by a classical
222 solution $\tilde{u}(t, x)$ of the regularized problem (2.1), where \tilde{f} is a smooth approximation of f
223 and $\tilde{u}(t, \cdot)$ has the same initial condition as $u(t, \cdot)$. We then aim to show that any three

224 interfaces $\tilde{\xi}_1(t) \leq \tilde{\xi}_2(t) \leq \tilde{\xi}_3(t)$ of $\tilde{u}(t, \cdot)$ coalesce to a single interface within finite time.
 225 We address the second challenge by approximating $\tilde{\xi}_2(t)$ on a compact time interval by
 226 a sequence of smooth approximations $\tilde{\xi}_{2,n}(t)$. Thus, the mass functions (3.1) with $u(t, x)$
 227 replaced by $\tilde{u}(t, x)$ and $\xi_2(t)$ by $\tilde{\xi}_{2,n}(t)$ are differentiable with respect to t and we can
 228 obtain the desired inequalities, which then yield that the interfaces $\tilde{\xi}_1(t), \tilde{\xi}_2(t)$ and $\tilde{\xi}_3(t)$
 229 of $\tilde{u}(t, x)$ coalesce to a single interface before an explicit time $T > 0$, which is independent
 230 of the approximation function \tilde{f} .

231 The approximation of the flux f by a smooth function \tilde{f} introduces an additional
 232 difficulty. Even with control on the norm $\|u(t, \cdot) - \tilde{u}(t, \cdot)\|_{W^{1,\infty}}$ through Lemma 2.3, the
 233 fact that $\tilde{u}(t, \cdot)$ possesses a single interface is not sufficient to conclude that $u(t, \cdot)$ has a
 234 single interface because interfaces of $u(t, \cdot)$ might accumulate close to the single interface
 235 of $\tilde{u}(t, \cdot)$. We address this issue by bounding the derivative $\partial_x \tilde{u}(t, \cdot)$ at the interface away
 236 from 0, precluding the accumulation of multiple interfaces of $u(t, \cdot)$ close to the single
 237 interface of $\tilde{u}(t, \cdot)$.

238 We bound the derivative of $\tilde{u}(t, \cdot)$ away from 0 by considering a traveling shock-wave
 239 solution $\tilde{u}_{\text{tw}}(t, x) = \psi(x - ct)$ of (2.1), which propagates at some speed $c \in \mathbb{R}$ and connects
 240 asymptotic limits ψ_{\pm} of opposite signs satisfying $|\psi_{\pm}| < |\phi_{\pm}|$. Upon switching to a co-
 241 moving frame, we may without loss of generality assume that $c = 0$. We then show, with
 242 the same methods as before, that all interfaces of the difference $v(t, \cdot) = \tilde{u}(t, \cdot) - \psi$ converge
 243 to a single interface within finite time, see Figure 3.1. This then yields the desired lower
 244 bound on $\|\partial_x \tilde{u}(t, \cdot)\|_{L^\infty}$. Using that $\|u(t, \cdot) - \tilde{u}(t, \cdot)\|_{W^{1,\infty}}$ can be taken sufficiently small
 245 by taking a better approximation \tilde{f} of f if necessary, we thus conclude that the solution
 246 $u(t, \cdot)$ must have a single interface for $t > T$, since the same holds for the approximation
 247 $\tilde{u}(t, \cdot)$.

248 Before we proceed with the proof of Theorem 3.1, we first state the following tech-
 249 nical lemma, which establishes a suitable smooth approximation \tilde{f} of the flux function
 250 f in (1.1). Naturally, we require that \tilde{f} lies sufficiently close to f and its derivative is
 251 well-behaved. Moreover, we wish that the regularized problem (2.1) admits a traveling
 252 shock-wave solution connecting the asymptotic states ϕ_{\pm} , but also a traveling shock wave
 253 with asymptotic limits ψ_{\pm} of opposite signs lying in between ϕ_- and ϕ_+ , see also Fig-
 254 ure 3.1. Without loss of generality, we can restrict to the case $\phi_- < 0 < \phi_+$ and we
 255 may assume $f(\phi_+) = f(\phi_-)$ by replacing $f(u)$ by $f(u) + cu$, where c is given by the
 256 Rankine-Hugoniot condition (1.3).

Lemma 3.2. *Let f be locally Lipschitz continuous and let $\phi_{\pm} \in \mathbb{R}$ with $\phi_- < 0 < \phi_+$. Suppose that $f(\phi_+) = f(\phi_-)$ and the Gel'fand-Oleinik entropy condition*

$$f(z) - f(\phi_{\pm}) < 0, \quad (3.2)$$

257 *holds for all $z \in (\phi_-, \phi_+)$. Then, for each $\kappa \in (f(0), f(\phi_{\pm}))$, there exists a constant*
 258 *$R > 0$ such that for all $\delta \in (0, \kappa - f(0))$, there exist $\tilde{f} \in C^\infty(\mathbb{R})$ and $\psi_{\pm} \in (\phi_-, \phi_+)$ with*
 259 *$\psi_- < 0 < \psi_+$ such that the following assertions hold:*




FIGURE 3.1. Left: the approximate shock-wave solution $\tilde{u}(t, \cdot)$ of (2.1) with asymptotic limits ϕ_{\pm} , the traveling shock wave ψ with asymptotic limits ψ_{\pm} and the interfaces $\tilde{\xi}_1(t)$, $\tilde{\xi}_2(t)$ and $\tilde{\xi}_3(t)$ of the difference $v(t, \cdot) = \tilde{u}(t, \cdot) - \psi$. In the proof of Theorem 3.1 we bound the shaded areas above and below the graph of $\tilde{u}(t, \cdot)$ from below by the orange subareas. Right: the smooth approximation \tilde{f} of the flux function f , established in Lemma 3.2. One observes that the regularized problem (2.1) admits a standing shock-wave solution connecting the asymptotic limits ϕ_{\pm} and one connecting the asymptotic states ψ_{\pm} , where $\phi_- < \psi_- < 0 < \psi_+ < \phi_+$.

i) For all $z \in (\phi_-, \phi_+)$ we have

$$\tilde{f}(\phi_+) = \tilde{f}(\phi_-), \quad \tilde{f}'(\phi_{\pm}) \neq 0, \quad \tilde{f}(z) - \tilde{f}(\phi_{\pm}) < 0. \quad (3.3)$$

ii) For all $z \in (\psi_-, \psi_+)$ it holds

$$\tilde{f}(\psi_+) = \kappa = \tilde{f}(\psi_-), \quad \tilde{f}'(\psi_{\pm}) \neq 0, \quad \tilde{f}(z) - \tilde{f}(\psi_{\pm}) < 0. \quad (3.4)$$

iii) For all $z \in [\phi_-, \phi_+]$ we have

$$|f(z) - \tilde{f}(z)| < \delta, \quad |\tilde{f}'(z)| < R. \quad (3.5)$$

Proof. We first recall that, since f is locally Lipschitz continuous, Rademacher's theorem asserts that f is differentiable almost everywhere and its derivative f' is essentially bounded on each bounded interval. We denote

$$R_1 := \sup\{|f'(u)| : u \in [\phi_-, \phi_+]\}.$$

Take $\delta \in (0, \kappa - f(0))$. Let $\Phi: \mathbb{R} \rightarrow \mathbb{R}$ be a mollifier with $\|\Phi\|_1 = 1$, $\Phi(x) > 0$ for $x \in (\phi_-, \phi_+)$ and $\Phi(x) = 0$ for $x \in \mathbb{R} \setminus (\phi_-, \phi_+)$. Set $\Phi_{\eta}(x) = \Phi(x/\eta)/\eta$ for $\eta > 0$. The function $g: \mathbb{R} \rightarrow \mathbb{R}$ given by $g(x) = \min\{f(x) + \frac{\delta}{4}, f(\phi_-)\}$ is locally Lipschitz continuous. Moreover, it holds $|g'(x)| \leq |f'(x)|$ for each $x \in [\phi_-, \phi_+]$. Since g is continuous, it can be approximated by the sequence $g_{\eta} := \Phi_{\eta} * g$ of smooth functions. That is, there exists $\eta_0 > 0$ such that

$$|g_{\eta}(u) - g(u)| < \frac{\delta}{4}$$

for all $u \in [\phi_-, \phi_+]$ and $\eta \in (0, \eta_0)$. By construction we have $g_\eta(x) \leq f(\phi_\pm)$ for all $x \in \mathbb{R}$ and $\eta > 0$. In addition, since g is constant in a neighborhood of ϕ_\pm and it holds $g'_\eta = \Phi_\eta * g'$, there exists $\eta_1 \in (0, \eta_0)$ such that $g_\eta(\phi_\pm) = f(\phi_\pm)$ and $|g'_\eta(u)| \leq \|\Phi_\eta\|_1 R_1 = R_1$ for all $\eta \in (0, \eta_1]$. We conclude that $\tilde{g} = g_{\eta_1} - \delta\Phi/(4\|\Phi\|_\infty)$ is a smooth function which satisfies $\tilde{g}(z) < \tilde{g}(\phi_\pm) = f(\phi_\pm)$ for $z \in (\phi_-, \phi_+)$. Moreover, it holds

$$|\tilde{g}(u) - f(u)| \leq |g_{\eta_1}(u) - g(u)| + \frac{\delta}{4} + |g(u) - f(u)| < \frac{3\delta}{4},$$

and

$$|\tilde{g}'(u)| \leq R_1 + \frac{\delta\|\Phi'\|_\infty}{4\|\Phi\|_\infty},$$

260 for $u \in [\phi_-, \phi_+]$.

Since we have $\tilde{g}(0) < f(0) + \delta < \kappa < f(\phi_\pm) = \tilde{g}(\phi_\pm)$, the open set $\tilde{g}^{-1}[\{z \in \mathbb{R} : z < \kappa\}]$ must contain an interval (ψ_-, ψ_+) with $\phi_- < \psi_- < 0 < \psi_+ < \phi_+$ and $\tilde{g}(\psi_+) = \kappa = \tilde{g}(\psi_-)$. Hence, it holds $\tilde{g}(z) < \tilde{g}(\psi_\pm)$ for $z \in (\psi_-, \psi_+)$. Finally, set $d = \frac{1}{2} \min\{\phi_+ - \psi_+, \psi_+, -\psi_-, \psi_- - \phi_-\} > 0$ and let $\Psi: \mathbb{R} \rightarrow \mathbb{R}$ be an even, smooth cut-off function such that $\Psi(0) = 1$, $\|\Psi\|_\infty \leq 1$, $\Psi(x) > 0$ for all $x \in (-d, d)$ and $\Psi(x) = 0$ for all $x \in \mathbb{R} \setminus (-d, d)$. Recalling the properties of the function \tilde{g} , we conclude that for any $\kappa_\pm, \lambda_\pm \in [0, \delta/(4(\phi_+ - \phi_-))]$, the smooth function

$$\begin{aligned} \tilde{f}(x) = & \tilde{g}(x) - \kappa_-(x - \phi_-)\Psi(x - \phi_-) - \lambda_-(x - \psi_-)\Psi(x - \psi_-) + \lambda_+(x - \psi_+)\Psi(x - \psi_+) \\ & + \kappa_+(x - \phi_+)\Psi(x - \phi_+), \end{aligned}$$

satisfies (3.5), it holds

$$\tilde{f}(\phi_+) = \tilde{f}(\phi_-), \quad \tilde{f}'(\phi_\pm) = \tilde{g}'(\phi_\pm) \pm \kappa_\pm, \quad \tilde{f}(z) - \tilde{f}(\phi_\pm) < 0,$$

for all $z \in (\phi_-, \phi_+)$, and we have

$$\tilde{f}(\psi_+) = \kappa = \tilde{f}(\psi_-), \quad \tilde{f}'(\psi_\pm) = \tilde{g}'(\psi_\pm) \pm \lambda_\pm, \quad \tilde{f}(z) - \tilde{f}(\psi_\pm) < 0, \quad (3.6)$$

261 for all $z \in (\psi_-, \psi_+)$. Hence, choosing $\kappa_\pm, \lambda_\pm \in [0, \delta/(4(\phi_+ - \phi_-))]$ in such a way that
 262 $\tilde{f}'(\phi_\pm), \tilde{f}'(\psi_\pm) \neq 0$, we find that \tilde{f} satisfies (3.3), (3.4), and (3.5). \square

263 Having established a suitable approximation \tilde{f} of the flux function f , we now provide
 264 the proof of Theorem 3.1 following the outline sketched above.

Proof of Theorem 3.1. We consider the case $\phi_- < 0 < \phi_+$. The case $\phi_+ < 0 < \phi_-$ is handled analogously. Clearly, the zeros (including their multiplicities) of $u(t, \cdot)$ are the same as those of the translate $u(t, \cdot - ct)$ for any $t \geq 0$. Thus, upon replacing $f(u)$ by $f(u) + cu$ in (1.1), where c is given by (1.3), we may assume

$$f(\phi_+) = f(\phi_-),$$

so that (1.2) yields

$$f(z) - f(\phi_+) = f(z) - f(\phi_-) < 0,$$

for all $z \in (\phi_-, \phi_+)$. By continuity of f there exists $\eta > 0$ such that for all $z \in [\phi_-, \phi_- + \eta] \cup [\phi_+ - \eta, \phi_+]$ it holds

$$f(z) > \frac{f(\phi_\pm) + f(0)}{2}. \quad (3.7)$$

Note that since $f(0) < f(\phi_\pm)$, we must have $\eta < |\phi_\pm|$. Since u_0 is continuous and converges to $\phi_\pm \neq 0$ as $x \rightarrow \pm\infty$, the function $u_0 - \phi_+ + \eta$ possesses a largest root ξ_+ and $u_0 - \phi_- - \eta$ possesses a smallest root ξ_- . We set

$$T = \frac{\max \left\{ \int_{-\infty}^{\xi_+} (u_0(x) - \phi_-) dx, \int_{\xi_-}^{\infty} (\phi_+ - u_0(x)) dx \right\}}{f(\phi_+) - f(0)} > 0. \quad (3.8)$$

We argue by contradiction and assume that there exists $\tau > T$ such that $u(\tau, \cdot)$ has at least two distinct zeros. Then, since u is continuously differentiable, there must exist a zero x_0 of $u(\tau, \cdot)$ with $u_x(\tau, x_0) \leq 0$. Fix $\kappa > f(0)$ such that

$$(\kappa - f(0))\tau < (f(\phi_-) - f(0))(\tau - T), \quad \kappa < \frac{f(0) + f(\phi_\pm)}{2}. \quad (3.9)$$

Denote by $L > 0$ the Lipschitz constant of f on $[\phi_-, \phi_+]$ and let $R > 0$ be the constant from Lemma 3.2 (which depends on κ). Fix $\varepsilon > 0$ such that

$$(L + 2)\varepsilon < \kappa - f(0), \quad (R + 1)\varepsilon < \kappa - f(0), \quad \varepsilon < \min\{M_\eta, m_\eta\}. \quad (3.10)$$

Finally, let $\delta_0 > 0$ be the constant from Lemma 2.3 (which depends on $R, \tau, \varepsilon > 0$) and take $\delta > 0$ such that

$$\begin{aligned} \delta < \min\{\delta_0, \varepsilon, \kappa - f(0)\}, \quad \delta < \frac{f(0) + f(\phi_\pm)}{2} - \kappa, \\ \delta\tau < (f(\phi_-) - f(0))(\tau - T) - (\kappa - f(0))\tau, \end{aligned} \quad (3.11)$$

265 which is possible by (3.9).

By Lemma 3.2 there exist $\tilde{f} \in C^\infty(\mathbb{R})$ and $\psi_\pm \in (\phi_-, \phi_+)$ with $\psi_- < 0 < \psi_+$ satisfying (3.3), (3.4), and (3.5). Lemma 2.3 then yields a global classical solution (2.3) of (2.1) with initial condition $\tilde{u}(0, \cdot) = u_0$ satisfying (2.5). Then, it must hold

$$\tilde{u}_x(\tau, x_0) \leq \varepsilon, \quad |\tilde{u}(\tau, x_0)| \leq \varepsilon. \quad (3.12)$$

On the other hand, the mean value theorem implies

$$\kappa - \tilde{f}(0) = \tilde{f}(\psi_\pm) - \tilde{f}(0) \leq R|\psi_\pm|.$$

Combining the latter with (3.5), and (3.10) yields

$$|\psi_\pm| \geq \frac{\kappa - f(0) - \delta}{R} \geq \frac{\kappa - f(0) - \varepsilon}{R} > \varepsilon. \quad (3.13)$$

On the other hand (3.4), (3.7), and (3.9) imply

$$\phi_- + \eta < \psi_- < 0 < \psi_+ < \phi_+ - \eta. \quad (3.14)$$

By (3.3) and (3.4) there exist heteroclinic solutions $\phi(x)$ and $\psi(x)$ of the profile equations

$$0 = \phi_\xi + \tilde{f}(\phi) - \tilde{f}(\phi_\pm), \quad 0 = \psi_\xi + \tilde{f}(\psi) - \tilde{f}(\psi_\pm), \quad (3.15)$$

266 respectively, converging exponentially to the asymptotic limits ϕ_\pm and ψ_\pm , respectively,
 267 as $x \rightarrow \pm\infty$. Since $u_0 - \phi_\pm$ is L^1 -integrable on \mathbb{R}_\pm , so is $u_0 - \phi$. Therefore, Lemma 2.4
 268 yields that $\tilde{u}(t, \cdot) - \phi$ is L^1 -integrable for all $t \geq 0$. We conclude that $\tilde{u}(t, \cdot) - \phi_\pm$ is
 269 L^1 -integrable on \mathbb{R}_\pm for all $t \geq 0$.

270 Using (3.12) and (3.13), and the fact that $\psi(x)$ is strictly monotone and converges to
 271 ψ_\pm as $x \rightarrow \pm\infty$, there must exist a translate $x_1 \in \mathbb{R}$ such that the point $(x_0, \tilde{u}(\tau, x_0))$ lies
 272 on the graph of $\psi(\cdot - x_1)$. Our aim is to show that the difference $v(t, \cdot) = \tilde{u}(t, \cdot) - \psi(\cdot - x_1)$
 273 has only a single zero at $t = \tau$, which must lie at x_0 . This then leads to a contradiction
 274 with (3.9), (3.11), (3.10), and (3.12) by our choice of constants κ, ε and δ .

Upon replacing the traveling shock wave ψ by its translate $\psi(\cdot - x_1)$, we may without loss of generality assume $x_1 = 0$. We observe that

$$v \in C([0, \infty), C_{\text{ub}}^1(\mathbb{R})) \cap C((0, \infty), C_{\text{ub}}^2(\mathbb{R})) \cap C^1((0, \infty), C_{\text{ub}}(\mathbb{R})), \quad (3.16)$$

is a global classical solution of the equation

$$v_t = v_{xx} + \tilde{f}(v + \psi)_x - \tilde{f}(\psi)_x. \quad (3.17)$$

We can apply the Sturm theorem, [1, Theorem B], upon recasting (3.17) as the linear parabolic equation

$$v_t = v_{xx} + b(t, x)v_x + a(t, x)v, \quad (3.18)$$

with

$$b(t, x) = \tilde{f}'(v(t, x) + \psi(t, x)), \quad a(t, x) = \psi_x(t, x) \frac{\tilde{f}'(v(t, x) + \psi(x)) - \tilde{f}'(\psi(x))}{v(t, x)},$$

275 where we note that $a, b, b_x,$ and b_t are bounded on the strip $\mathbb{R} \times [0, s]$ for any $s > 0$
 276 by (3.16), and the fact that \tilde{f} and ψ and smooth. Applying [1, Theorem B] to (3.18)
 277 yields that, if it holds $v(t_0, \xi_0) = 0 = v_x(t_0, \xi_0)$ at some $(t_0, \xi_0) \in (0, \infty) \times \mathbb{R}$, then there
 278 exist $\theta \in (0, t_0)$ and a neighborhood $U \subset \mathbb{R}$ of ξ_0 such that for $t \in (t_0 - \theta, t_0)$, there are
 279 at least two zeros of $v(t, \cdot)$ in U and for $t \in (t_0, t_0 + \theta)$, there is at most one zero of $v(t, \cdot)$
 280 in U . Noting that $v(t, x)$ is continuously differentiable with respect to x and t , this leads
 281 to two important observations. First, no new zeros of $v(t, \cdot)$ can form dynamically over
 282 time. Second, multiple roots are isolated in $\mathbb{R} \times (0, \infty)$.

283 Now assume by contradiction that for all $t \in [0, \tau]$, there exist at least two zeros of
 284 $v(t, \cdot)$. A consequence of the above two observations, the regularity of $v(t, \cdot)$, and the fact
 285 that $v(t, \cdot)$ converges to $\phi_\pm - \psi_\pm$ at $\pm\infty$ with $\phi_- - \psi_- < 0 < \phi_+ - \psi_+$, is that there must
 286 be three functions $\tilde{\xi}_{1,2,3}: [0, T] \rightarrow \mathbb{R}$ which depend continuously on time such that it holds
 287 $\tilde{\xi}_1(t) < \tilde{\xi}_2(t) < \tilde{\xi}_3(t)$, $v(t, \tilde{\xi}_i(t)) = 0$ for $i = 1, 2, 3$, $v(t, x) > 0$ for all $x \in (\tilde{\xi}_1(t), \tilde{\xi}_2(t))$,
 288 $v(t, x) < 0$ for all $x \in (\tilde{\xi}_2(t), \tilde{\xi}_3(t))$, and $v_x(t, \tilde{\xi}_2(t)) \leq 0$ for all $t \in [0, T]$. We note that
 289 by (3.14), it must hold $\xi_- < \tilde{\xi}_2(0) < \xi_+$.

Take a sequence $\{\tilde{\xi}_{2,n}\}_n$ of smooth functions converging uniformly in $C([0, \tau])$ to $\tilde{\xi}_2$ as $n \rightarrow \infty$. Define the masses

$$\begin{aligned} M_{1,n}(t) &= \int_{-\infty}^{\tilde{\xi}_{2,n}(t)} (v(t, x) - \phi_- + \psi_-) dx \\ &= \int_{-\infty}^{\tilde{\xi}_{2,n}(t)} (\tilde{u}(t, x) - \phi_-) dx - \int_{-\infty}^{\tilde{\xi}_{2,n}(t)} (\psi(x) - \psi_-) dx, \\ M_1(t) &= \int_{-\infty}^{\tilde{\xi}_2(t)} (v(t, x) - \phi_- + \psi_-) dx = \int_{-\infty}^{\tilde{\xi}_2(t)} (\tilde{u}(t, x) - \phi_-) dx - \int_{-\infty}^{\tilde{\xi}_2(t)} (\psi(x) - \psi_-) dx, \end{aligned}$$

which are well-defined as $\tilde{u}(t, \cdot) - \phi_-$ and $\psi - \psi_-$ are L^1 -integrable on \mathbb{R}_- for all $t \geq 0$. Applying the Leibniz' rule, we find

$$\begin{aligned} M'_{1,n}(s) &= \tilde{\xi}'_{2,n}(s) \left(v(s, \tilde{\xi}_{2,n}(s)) - \phi_- + \psi_- \right) \\ &\quad + \int_{-\infty}^{\tilde{\xi}_{2,n}(s)} \left(v_{xx}(s, x) + \partial_x \left(\tilde{f}(v(s, x) + \psi(x)) - \tilde{f}(\psi(x)) \right) \right) dx \\ &= \partial_s \left(\tilde{\xi}_{2,n}(s) \left(v(s, \tilde{\xi}_{2,n}(s)) - \phi_- + \psi_- \right) \right) - \tilde{\xi}_{2,n}(s) \partial_s \left(v(s, \tilde{\xi}_{2,n}(s)) \right) + v_x(s, \tilde{\xi}_{2,n}(s)) \\ &\quad + \tilde{f}(v(s, \tilde{\xi}_{2,n}(s)) + \psi(\tilde{\xi}_{2,n}(s))) - \tilde{f}(\psi(\tilde{\xi}_{2,n}(s))) - \tilde{f}(\phi_-) + \tilde{f}(\psi_-), \end{aligned}$$

for $s \in (0, \tau]$. Integrating the latter from 0 to t we obtain

$$\begin{aligned} M_{1,n}(t) &= M_{1,n}(0) + \tilde{\xi}_{2,n}(t) \left(v(t, \tilde{\xi}_{2,n}(t)) - \phi_- + \psi_- \right) - \tilde{\xi}_{2,n}(0) \left(v(0, \tilde{\xi}_{2,n}(0)) - \phi_- + \psi_- \right) \\ &\quad + \int_0^t \left(\tilde{\xi}_{2,n}(s) \partial_s \left[v(s, \tilde{\xi}_{2,n}(s)) \right] + v_x(s, \tilde{\xi}_{2,n}(s)) \right) ds + \left(\tilde{f}(\psi_-) - \tilde{f}(\phi_-) \right) t \\ &\quad + \int_0^t \left(\tilde{f}(v(s, \tilde{\xi}_{2,n}(s)) + \psi(\tilde{\xi}_{2,n}(s))) - \tilde{f}(\psi(\tilde{\xi}_{2,n}(s))) \right) ds, \end{aligned}$$

for $t \in (0, \tau]$. Taking the limit $n \rightarrow \infty$, while recalling the regularity (3.16) of $v(t, \cdot)$ and the fact that $v_x(\tilde{\xi}_2(s), s) \leq 0$ for all $s \in [0, T]$, we arrive at

$$\begin{aligned} M_1(t) &= M_1(0) - (\tilde{\xi}_2(t) - \tilde{\xi}_2(0)) (\phi_- - \psi_-) + \int_0^t v_x(s, \tilde{\xi}_2(s)) ds + \left(\tilde{f}(\psi_-) - \tilde{f}(\phi_-) \right) t \\ &\leq M_1(0) + (\tilde{\xi}_2(t) - \tilde{\xi}_2(0)) (\psi_- - \phi_-) + \left(\tilde{f}(\psi_-) - \tilde{f}(\phi_-) \right) t, \end{aligned}$$

implying

$$\begin{aligned} \int_{-\infty}^{\tilde{\xi}_2(t)} (\tilde{u}(t, x) - \phi_-) dx &\leq \int_{-\infty}^{\tilde{\xi}_2(0)} (u_0(x) - \phi_-) dx + \int_{\tilde{\xi}_2(0)}^{\tilde{\xi}_2(t)} (\psi(x) - \phi_-) dx \\ &\quad + \left(\tilde{f}(\psi_-) - \tilde{f}(\phi_-) \right) t, \end{aligned}$$

for $t \in [0, \tau]$. On the other hand, since $\tilde{u}(t, \cdot) - \phi_-$ is nonnegative for all $t \geq 0$ by (2.5), it holds

$$\int_{\tilde{\xi}_1(t)}^{\tilde{\xi}_2(t)} (\psi(x) - \phi_-) dx \leq \int_{-\infty}^{\tilde{\xi}_2(t)} (\tilde{u}(t, x) - \phi_-) dx, \quad (3.19)$$

cf. Figure 3.1. Combining the latter two inequalities, while using $\tilde{\xi}_2(0) < \xi_+$, we obtain

$$\begin{aligned} \int_{\tilde{\xi}_1(t)}^{\tilde{\xi}_2(0)} (\psi(x) - \phi_-) dx &\leq \int_{-\infty}^{\tilde{\xi}_2(0)} (u_0(x) - \phi_-) dx + \left(\tilde{f}(\psi_-) - \tilde{f}(\phi_-) \right) t \\ &\leq \int_{-\infty}^{\xi_+} (u_0(x) - \phi_-) dx + (f(0) - f(\phi_-)) t + (\kappa - f(0)) t + \delta t. \end{aligned}$$

Inserting $t = \tau$ in the latter, applying (3.11), and recalling (3.8), we arrive at

$$\int_{\tilde{\xi}_1(t)}^{\tilde{\xi}_2(0)} (\psi(x) - \phi_-) dx \leq (f(0) - f(\phi_-)) (\tau - T) + (\kappa - f(0)) \tau + \delta \tau < 0.$$

290 yielding $\tilde{\xi}_2(0) \leq \tilde{\xi}_1(\tau) < \tilde{\xi}_2(\tau)$, since we have $\psi(x) - \phi_- \geq \psi_- - \phi_- > 0$ for all $x \in \mathbb{R}$.

Similarly, we establish

$$\int_{\tilde{\xi}_2(0)}^{\tilde{\xi}_3(t)} (\phi_+ - \psi(x)) dx \leq \int_{\tilde{\xi}_2(0)}^{\infty} (\phi_+ - u_0(x)) dx + \left(\tilde{f}(\psi_+) - \tilde{f}(\phi_+) \right) t,$$

yielding

$$\int_{\tilde{\xi}_2(0)}^{\tilde{\xi}_3(t)} (\phi_+ - \psi(x)) dx \leq (f(0) - f(\phi_+)) (\tau - T) + (\kappa - f(0)) \tau + \delta \tau < 0,$$

and thus, $\tilde{\xi}_2(\tau) < \tilde{\xi}_3(\tau) \leq \tilde{\xi}_2(0)$, which contradicts $\tilde{\xi}_2(0) < \tilde{\xi}_2(\tau)$. Hence, there must exist a $t \in [0, \tau]$ such that $v(t, \cdot)$ has only a single zero. Recalling that the number of zeros is non-increasing, we conclude that $v(\tau)$ has a single zero, which must be x_0 . Since $v(t, \cdot)$ converges to $\phi_{\pm} - \psi_{\pm}$ as $x \rightarrow \pm\infty$ and we have $\phi_- < \psi_- < 0 < \psi_+ < \phi_+$, it must hold $\tilde{u}_x(\tau, x_0) - \psi'(x_0) = v_x(\tau, x_0) \geq 0$. On the other hand, using that ψ solves (3.15) and $0 = v(\tau, x_0) = \tilde{u}(\tau, x_0) - \psi(x_0)$, while recalling (3.5) and (3.12), we infer

$$\begin{aligned} \varepsilon &\geq \tilde{u}_x(\tau, x_0) \geq \psi'(x_0) = -\tilde{f}(\psi(x_0)) + \tilde{f}(\psi_{\pm}) = -\tilde{f}(\tilde{u}(\tau, x_0)) + \kappa \\ &= -\tilde{f}(\tilde{u}(\tau, x_0)) + f(\tilde{u}(\tau, x_0)) - f(\tilde{u}(\tau, x_0)) + f(0) + \kappa - f(0) \\ &\geq -\delta - L\varepsilon + \kappa - f(0). \end{aligned}$$

Combining the latter with (3.11) yields

$$(L + 2)\varepsilon \geq \kappa - f(0),$$

291 which contradicts (3.10). We conclude that for each $t > T$, the function $u(t, \cdot)$ possesses
292 at most one zero. \square

293 **Remark 3.3.** We note that the proof of Theorem 3.1 provides an explicit upper bound
 294 T , given by (3.8), on the time at which all interfaces of the solution $u(t, \cdot)$ of (1.1) have
 295 collapsed to a single interface. The upper bound (3.8) only depends on the flux function
 296 f and the initial condition u_0 .

Remark 3.4. We expect that it might be possible to lift the assumption that $u_0(x) \in$
 $[\min\{\phi_-, \phi_-\}, \max\{\phi_-, \phi_+\}]$ for all $x \in \mathbb{R}$ in Theorem 3.1 by bounding $u_0(x)$ from below
 by a smooth function $u_-(x)$ and from above by a smooth function $u_+(x)$ satisfying

$$\lim_{x \rightarrow \pm\infty} u_-(x) = \min\{\phi_-, \phi_+\} \quad \text{and} \quad \lim_{x \rightarrow \pm\infty} u_+(x) = \max\{\phi_-, \phi_+\}.$$

297 It has been established in [4] that the solutions $\tilde{u}_\pm(t, \cdot)$ of the regularized problem (2.1)
 298 with initial conditions $\tilde{u}_\pm(0, \cdot) = u_\pm$ converge in L^1 - and L^∞ -norm to their asymptotic
 299 limits as $t \rightarrow \infty$. So, by the comparison principle, the area of $\tilde{u}(t, \cdot)$ under $\min\{\phi_-, \phi_+\}$
 300 or above $\max\{\phi_-, \phi_+\}$ converges to 0 as $t \rightarrow \infty$. We expect that using similar techniques
 301 as in the proof of Theorem 3.5, one can obtain decay estimates on this area, which are
 302 independent of the approximation \tilde{f} of the flux function f . One would then hope to find
 303 an explicit time $T_1 > 0$, only depending on f and the initial condition u_0 , such that for
 304 $t > T_1$ this area is so small that the estimate (3.19) is still valid and one can proceed as
 305 in the proof of Theorem 3.1. We decided to refrain from providing this exposition, since
 306 it merely introduces additional technicalities obscuring the main ideas of the proof.

307 **3.2. Solutions with initial data of class II.** We prove the finite-time extinction of all
 308 interfaces of solutions with initial data of class II. That is, we consider a solution $u(t, x)$
 309 of (1.1) with initial condition $u(0, x) = u_0(x)$, which converges to nonzero asymptotic
 310 limits ϕ_\pm as $x \rightarrow \pm\infty$ that have the same sign. By approximating the solution $u(t, x)$ by a
 311 solution $\tilde{u}(t, x)$ to the regularized problem (2.1) with smooth flux function \tilde{f} and bounding
 312 the initial condition u_0 from below or above, it suffices by the comparison principle of [14,
 313 17] to prove the statement for a solution $\tilde{v}(t, \cdot)$ of the regularized problem (2.1) which
 314 possesses the same non-zero asymptotic limit ϕ_0 at $\pm\infty$, see Figure 3.2. We show that all
 315 interfaces of $\tilde{v}(t, \cdot)$ go extinct within finite time by deriving an energy inequality for the
 316 difference $\tilde{v}(t, \cdot) - \phi_0$. The energy estimate relies on the Gagliardo-Nirenberg inequality
 317 and the conservation of mass.

318 **Theorem 3.5.** Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be locally Lipschitz continuous and $u_0 \in C_{\text{ub}}^1(\mathbb{R})$. Suppose
 319 that $u_0(x)$ converges to nonzero asymptotic limits ϕ_\pm as $x \rightarrow \pm\infty$ such that ϕ_+ and
 320 ϕ_- have the same sign. Let $u \in C([0, \infty), C_{\text{ub}}(\mathbb{R}))$ be the global mild solution of (1.1),
 321 established in Lemma 2.2. Then, there exists a time $T > 0$ such that for all $t > T$, the
 322 solution $u(t, \cdot)$ possesses no zeros.

Proof. Throughout the proof, $C > 0$ denotes the constant appearing in the Gagliardo-Nirenberg interpolation inequality

$$\|g\|_\infty \leq C \|g'\|_2^{\frac{2}{3}} \|g\|_1^{\frac{1}{3}}, \quad (3.20)$$

323 which holds for all $g \in L^1(\mathbb{R}) \cap H^1(\mathbb{R})$.

We consider the case $0 < \phi_- \leq \phi_+$. The cases $0 < \phi_+ \leq \phi_-$, $\phi_- \leq \phi_+ < 0$ and $\phi_+ \leq \phi_- < 0$ are handled analogously. Take any $v_0 \in C_{\text{ub}}^1(\mathbb{R})$ such that $v_0 - \frac{2}{3}\phi_-$ is L^1 -integrable, not identically zero, and nonpositive and it holds $v_0(x) \leq u_0(x)$ for all $x \in \mathbb{R}$. Set

$$T = \frac{2\phi_-^3}{9C^3 \left\| v_0 - \frac{2}{3}\phi_- \right\|_2^2 \left\| v_0 - \frac{2}{3}\phi_- \right\|_1} > 0. \quad (3.21)$$

Let $\tau > T$. By Lemma 2.2 there exists $\tilde{f} \in \mathbb{C}^\infty(\mathbb{R})$ such that the global classical solution $\tilde{u}(t, \cdot)$ of the regularized problem (2.1) with initial condition $\tilde{u}(0, \cdot) = u_0$ satisfies (2.3) and

$$\|u(\tau, \cdot) - \tilde{u}(\tau, \cdot)\|_\infty < \frac{1}{3}\phi_-. \quad (3.22)$$

Let

$$\tilde{v} \in C([0, \infty), C_{\text{ub}}^1(\mathbb{R})) \cap C((0, \infty), C_{\text{ub}}^2(\mathbb{R})) \cap C^1((0, \infty), C_{\text{ub}}(\mathbb{R})),$$

be the solution of (2.1) with initial condition $\tilde{v}(0, \cdot) = v_0$, cf. Lemma 2.1. By the comparison principle, cf. [14, 17], it holds

$$\tilde{v}(t, x) \leq \tilde{u}(t, x), \quad \tilde{v}(t, x) \leq \frac{2}{3}\phi_-, \quad (3.23)$$

324 for all $t \geq 0$ and $x \in \mathbb{R}$. Our aim is to show that we have $\tilde{v}(\tau, x) \geq \frac{1}{3}\phi_-$ for all $x \in \mathbb{R}$,
 325 which together with (3.22) and (3.23) yields the desired result that $u(\tau, \cdot)$ does not posses
 326 any zeros, cf. Figure 3.2.




FIGURE 3.2. Left: the shock wave $u(t, \cdot)$ and its approximation $\tilde{u}(t, \cdot)$ from the proof of Theorem 3.1 at time $t = \tau$. The shock wave $u(\tau, \cdot)$ possesses the asymptotic limits ϕ_\pm at $\pm\infty$ and has an interface at x_0 . The translate $\psi(\cdot - x_1)$ of the traveling shock wave ψ , connecting the asymptotic states ψ_\pm , passes through the point $(x_0, \tilde{u}(\tau, x_0))$. Right: the approximate solution $\tilde{u}(t, \cdot)$ connecting the asymptotic end states ϕ_\pm and its subsolution $\tilde{v}(t, \cdot)$ possessing the asymptotic limit $\frac{2}{3}\phi_-$ at $\pm\infty$. In the proof of Theorem 3.5 we approximate the energy of $\tilde{v}(t, \cdot) - \frac{2}{3}\phi_-$ at a point $t = \tau$ from below by $(\xi_2 - \xi_1)\frac{1}{9}\phi_-^2$.

We argue by contradiction and assume that there exist $\xi_1, \xi_2 \in \mathbb{R}$ with $\xi_1 < \xi_2$ such that $\tilde{v}(\tau, \xi_1) = \frac{1}{3}\phi_- = \tilde{v}(\tau, \xi_2)$. First, we observe that (τ, ξ_1) is a root of $z(t, x) = \tilde{v}(t, x) - \frac{1}{3}\phi_-$, which satisfies the linear equation

$$z_t = z_{xx} + b(t, x)z_x, \quad (3.24)$$

where the spatial and temporal derivative of $b(t, x) = \tilde{f}'(\tilde{v}(t, x))$ are bounded on the strip $\mathbb{R} \times [0, s]$ for any $s > 0$. Applying the Sturm Theorem [1, Theorem B] to (3.24) yields that $z(t, \cdot)$ must have a zero for all $t \in [0, \tau]$. That is, it holds

$$\|\tilde{v}(t, \cdot) - \frac{2}{3}\phi_-\|_\infty \geq \frac{1}{3}\phi_-, \quad (3.25)$$

327 for all $t \in [0, \tau]$.

Next, we observe that the mass

$$M(t) = \int_{\mathbb{R}} (\tilde{v}(t, x) - \frac{2}{3}\phi_-) dx, \quad t \geq 0,$$

is conserved. Indeed, it holds

$$M'(t) = \int_{\mathbb{R}} (\tilde{v}_{xx}(t, x) + \partial_x(\tilde{f}(\tilde{v}(t, x)))) dx = 0,$$

and thus, we have $M(t) = M(0)$ for all $t \geq 0$. Second, we establish an estimate for the energy

$$E(t) = \|\tilde{v}(t, \cdot) - \frac{2}{3}\phi_-\|_2^2.$$

We compute using integration by parts

$$\begin{aligned} E'(t) &= 2 \int_{\mathbb{R}} (\tilde{v}(t, x) - \frac{2}{3}\phi_-) (\tilde{v}_{xx}(t, x) + \partial_x(\tilde{f}(\tilde{v}(t, x)))) dx \\ &= -2 \int_{\mathbb{R}} \tilde{v}_x(t, x) (\tilde{v}_x(t, x) + \tilde{f}'(\tilde{v}(t, x))) dx \\ &= -2\|\tilde{v}_x(t, \cdot)\|_2^2, \end{aligned}$$

for $t \geq 0$. Therefore, using the Gagliardo-Nirenberg inequality (3.20), the bound (3.23) and the fact that the (nonzero) mass $M(t)$ is conserved, we obtain the energy estimate

$$E'(t) \leq -\frac{2}{C^3|M(t)|} \|\tilde{v}(t, \cdot) - \frac{2}{3}\phi_-\|_\infty^3 = -\frac{2}{C^3|M(0)|} \|\tilde{v}(t, \cdot) - \frac{2}{3}\phi_-\|_\infty^3,$$

for $t \geq 0$. Integrating the latter from 0 to τ , while using (3.25) and $\tau > T$, we obtain

$$\begin{aligned} (\xi_2 - \xi_1) \frac{\phi_-^2}{9} &\leq E(\tau) \leq E(0) - \frac{2}{C^3|M(0)|} \int_0^\tau \|\tilde{v}(t, \cdot) - \frac{2}{3}\phi_-\|_\infty^3 dt \\ &\leq E(0) - \frac{2\phi_-^3}{9C^3|M(0)|} \tau = E(0) \left(1 - \frac{\tau}{T}\right) < 0, \end{aligned}$$

328 which contradicts $\xi_1 < \xi_2$, see Figure 3.2. Therefore, $\tilde{v}(\tau, \cdot) - \frac{1}{3}\phi_-$ can possess at most
 329 one single zero, which together with estimates (3.22) and (3.23) and the fact that $\tilde{v}(\tau, x)$
 330 converges to $\frac{2}{3}\phi_-$ as $x \rightarrow \pm\infty$, implies that $u(\tau, \cdot)$ cannot have any zeros. \square

Remark 3.6. *Assume that the initial condition u_0 in Theorem 3.5 possesses an interface and it holds $0 < \phi_- \leq \phi_+$. By mollifying the compactly supported, nonpositive, nonzero function $u_1(x) = \min\{\frac{2}{3}\phi_-, u_0(x)\} - \frac{2}{3}\phi_-$, one readily finds a sequence $\{z_n\}_n$ of nonpositive, nonzero, smooth, and compactly supported functions such that z_n converges in $L^p(\mathbb{R})$ to u_1 as $n \rightarrow \infty$ for $p = 1, 2$. Thus, $w_n = z_n + \frac{2}{3}\phi_-$ is a smooth function such that $w_n - \frac{2}{3}\phi_-$ is L^1 -integrable, not identically zero, and nonpositive such that $w_n(x) \leq u_0(x)$ for all $n \in \mathbb{N}$. Hence, w_n satisfies the criteria for the function v_0 in the proof of Theorem 3.5 for any $n \in \mathbb{N}$. That is, we find that the upper bound (3.21) on the time at which all interfaces of the solution $u(t, \cdot)$ have gone extinct, could be taken equal to*

$$T = \frac{2\phi_-^3}{9C^3\|u_1\|_2^2\|u_1\|_1}.$$

331 *We stress that T only depends on the initial condition u_0 of the solution $u(t, \cdot)$ and the*
 332 *positive constant C from the Gagliardo-Nirenberg inequality (3.20).*

333 **3.3. Solutions with initial data of class III.** In Theorem 3.1, we proved finite-time
 334 coalescence of interfaces for shock waves converging to asymptotic limits of opposite signs.
 335 This prompts the question of whether anti-shock waves converging to asymptotic limits of
 336 opposite signs also exhibit finite-time coalescence of interfaces. One readily observes that
 337 the proof of Theorem 3.1 strongly relies on the Gel'fand-Oleinik entropy inequality (1.2) to
 338 bound the mass. It cannot be expected that the same strategy applies to the case of anti-
 339 shock waves that violate (1.2). Therefore, the question of whether finite-time coalescence
 340 of interfaces can be established for solutions with initial data of class III remains open.

341 Nevertheless, we can study the interface dynamics of solutions with initial data of class
 342 III in the framework of the modular Burgers' equation

$$343 \quad u_t = u_{xx} + |u|_x, \tag{3.26}$$

344 which corresponds to the scalar viscous conservation law (1.1) with the modular flux func-
 345 tion $f(u) = |u|$. Our upcoming analysis establishes finite-time coalescence of interfaces
 346 for anti-shock waves converging to asymptotic limits $\mp\phi_*$ as $x \rightarrow \pm\infty$ with $\phi_* > 0$. We
 347 make the following assumption on the regularity of solutions to the modular Burgers'
 348 equation (3.26).

349 **Assumption 3.7.** *For every $u_0 \in C_{\text{ub}}^1(\mathbb{R})$ converging to nonzero asymptotic limits u_{\pm} at*
 350 *$\pm\infty$, the global mild solution $u \in C([0, \infty), C_{\text{ub}}(\mathbb{R}))$ of (3.26), established in Lemma 2.2,*
 351 *with initial condition $u(0, \cdot) = u_0$ satisfies $u \in C^1((0, \infty) \times \mathbb{R}, \mathbb{R})$ such that $u(t, \cdot)$, $t \geq 0$*
 352 *is piecewise C^2 with the finite jump condition*

$$353 \quad u_{xx}(t, \xi(t)^+) - u_{xx}(t, \xi(t)^-) = -2|u_x(t, \xi(t))|, \tag{3.27}$$

354 *across any interface $x = \xi(t) \in \mathbb{R}$.*

355 Assumption 3.7 was proven in [9] for the class of solutions to (3.26) with a single
 356 interface in a local neighborhood of a traveling shock wave. In a more general setting, the
 357 validity of Assumption 3.7 is an open question.

358 We expect that Assumption 3.7 can be proven in a general case by using approximation
 359 by solutions of the regularized equation as in Theorems 3.1 and 3.5. However, since our
 360 main goal is to illustrate the finite-time coalescence of interfaces of solutions of (1.1) with
 361 initial data of class III rather than proving a general well-posedness result for piecewise
 362 smooth flux functions, we refrain from doing so.

363 The following lemma establishes that the odd parity of initial data is preserved in the
 364 time evolution of the modular Burgers' equation (3.26).

365 **Lemma 3.8.** *Let $u_0 \in C_{\text{ub}}^1(\mathbb{R})$ satisfy $u_0(-x) = -u_0(x)$ for every $x \in \mathbb{R}$. Then, the mild*
 366 *solution $u \in C([0, \infty), C_{\text{ub}}(\mathbb{R}))$ of (3.26), established in Lemma 2.2, satisfies $u(t, -x) =$*
 367 *$-u(t, x)$ for every $t \geq 0$ and $x \in \mathbb{R}$.*

Proof. First, observe that, if $z \in C_{\text{ub}}(\mathbb{R})$ is odd, then

$$e^{\partial_x^2 t} z = \int_{\mathbb{R}} \frac{e^{-\frac{y^2}{4t}}}{\sqrt{4\pi t}} z(x-y) dy$$

is also odd, which follows by the substitution $y \mapsto -y$. Now the mild solution $u(t, \cdot)$ of (3.26) is given by

$$u(t, \cdot) = e^{\partial_x^2 t} u_0 + \partial_x \int_0^t e^{\partial_x^2(t-s)} |u(s, \cdot)| ds,$$

for $t \geq 0$. Since u_0 is odd, so is $e^{\partial_x^2 t} u_0$. Hence, using again the substitution $y \mapsto y$, we obtain

$$u(t, \cdot) + u(t, -\cdot) = \partial_x \int_0^t e^{\partial_x^2(t-s)} (|u(s, \cdot)| - |u(s, -\cdot)|) ds,$$

for $t \geq 0$. Taking norms in the latter and recalling the well-known fact that there exists a constant $C > 0$ such that

$$\|\partial_x e^{\partial_x^2 t} v\|_{L^\infty} \leq Ct^{-1/2} \|v\|_{L^\infty}$$

for $t > 0$ and $v \in L^\infty(\mathbb{R})$, yields

$$\begin{aligned} \|u(t, \cdot) + u(t, -\cdot)\|_{L^\infty} &\leq C \int_0^t \frac{1}{\sqrt{t-s}} \| |u(s, \cdot)| - |u(s, -\cdot)| \|_{L^\infty} ds \\ &\leq C \int_0^t \frac{1}{\sqrt{t-s}} \|u(s, \cdot) + u(s, -\cdot)\|_{L^\infty} ds, \end{aligned}$$

for $t \geq 0$. Therefore, Grönwall's inequality, cf. [10, Lemma 7.0.3], implies that

$$\|u(t, \cdot) + u(t, -\cdot)\|_{L^\infty} = 0$$

368 for all $t \geq 0$, which finishes the proof. □

369 The main result of this section is the following theorem.

Theorem 3.9. *Suppose Assumption 3.7 holds. Take $\phi_* > 0$ and $x_{0,1} \in \mathbb{R}$ with $0 < x_1 - x_0 < \frac{1}{6}$. Let $\phi: \mathbb{R} \rightarrow \mathbb{R}$ be the odd function given by*

$$\phi(x) = \phi_*(e^{-x} - 1),$$

for $x \geq 0$. Consider $u_0 \in C_{\text{ub}}^1(\mathbb{R})$ satisfying

$$\phi(x - x_0) \leq u_0(x) \leq \phi(x - x_1), \quad (3.28)$$

370 for all $x \in \mathbb{R}$. Let $u \in C([0, \infty), C_{\text{ub}}(\mathbb{R}))$ be the mild solution of (3.26) established in
 371 Lemma 2.2. Then, $u(t, \cdot)$ cannot possess two consecutive simple zeros $\xi_1(t), \xi_2(t)$ that exist
 372 for all $t \geq 0$.

Proof. Our analysis relies on comparison with an explicit reference solution $u_{\text{ref}}(t, x)$ of (3.26) with odd initial condition $u_{\text{ref}}(0, \cdot) = \phi \in C_{\text{ub}}^1(\mathbb{R})$. By Lemma 3.8, the solution $u_{\text{ref}} \in C([0, \infty), C_{\text{ub}}(\mathbb{R}))$ is spatially odd. It satisfies the following diffusion-advection boundary-value problem:

$$\begin{cases} u_t = u_{xx} - u_x, & t > 0, \quad x > 0, \\ u(t, 0) = 0, & t \geq 0, \\ u(0, x) = \phi(x), & x \geq 0, \end{cases}$$

whose solution is explicitly given by

$$u_{\text{ref}}(t, x) = \int_0^\infty G(t, x, y) \phi(y) dy,$$

for $t \geq 0$ and $x \in \mathbb{R}$, where $G(t, x, y)$ is the Green's function used in [9]:

$$G(t, x, y) = \frac{1}{\sqrt{4\pi t}} \left(e^{-\frac{(x-y-t)^2}{4t}} - e^{-y} e^{-\frac{(x+y-t)^2}{4t}} \right).$$

Evaluating the integral we find

$$\begin{aligned} u_{\text{ref}}(t, x) = & \frac{\phi_*}{2} \left(e^x \operatorname{erfc} \left(\frac{t+x}{2\sqrt{t}} \right) - \operatorname{erfc} \left(\frac{t-x}{2\sqrt{t}} \right) \right. \\ & \left. - e^{2t-x} \left(\operatorname{erfc} \left(\frac{x-3t}{2\sqrt{t}} \right) + e^{3x} \operatorname{erfc} \left(\frac{3t+x}{2\sqrt{t}} \right) - 2 \right) \right), \end{aligned}$$

373 for $t \geq 0$ and $x \in \mathbb{R}$.

By the comparison principle, cf. [3, Corollary 3.1], and (3.28) it holds

$$u_-(t, x) \leq u(t, x) \leq u_+(t, x)$$

374 for $x \in \mathbb{R}$ and $t \geq 0$, where $u_-(t, x) = u_{\text{ref}}(t, x - x_0)$ and $u_+(t, x) = u_{\text{ref}}(t, x - x_1)$ are
 375 translates of the reference solution $u_{\text{ref}}(t, x)$ of (3.26), see Figure 3.3. Note that $u_-(t, \cdot)$
 376 and $u_+(t, \cdot)$ possess an odd symmetry with respect to the points $x = x_0$ and $x = x_1$,
 377 respectively. In particular, it holds $u_-(t, x_0) = 0 = u_+(t, x_1)$.

378 We argue by contradiction and assume that $u(t, x)$ possesses zeros $\xi_1(t), \xi_2(t), \xi_3(t)$ for
 379 all $t \geq 0$ such that $\xi_1(t) < \xi_2(t) < \xi_3(t)$, $u(t, x) < 0$ for $x \in (\xi_1(t), \xi_2(t))$, $u(t, x) > 0$ for all
 380 $x \in (\xi_2(t), \xi_3(t))$, and $u_x(t, \xi_2(t)) > 0$ for all $t \geq 0$. Since $u_\pm(t, \cdot)$ are monotone, it holds

381 $\xi_i(t) \in (x_0, x_1)$ for all $t \geq 0$ and $i = 1, 2, 3$, see Figure 3.3. By translational invariance, we
 382 may assume without loss of generality that $x_0 = 0$.

As in the proof of Theorem 3.1, we derive differential inequalities for the masses

$$M_1(t) = \int_{-\infty}^{\xi_2(t)} (\phi_* - u(t, x)) dx, \quad M_2(t) = \int_{\xi_2(t)}^{\infty} (u(t, x) + \phi_*) dx.$$

383 However, in contrast to the proof of Theorem 3.1, we cannot employ the Gel'fand-Oleinik
 384 entropy inequality to bound $M_1(t)$ and $M_2(t)$. Instead, we use explicit expressions of the
 385 reference solutions $u_{\pm}(t, \cdot)$ to bound $M_1(0)$ and $M_2(0)$ from above and $M_1(t)$ and $M_2(t)$
 386 from below.

Recalling $u_x(\xi_2(t), 0) > 0$, the implicit function theorem implies that $\xi_2(t)$ is differentiable with respect to t . We apply Leibniz' rule to compute

$$\begin{aligned} M_2'(t) &= -\xi_2'(t)\phi_* + \int_{\xi_2(t)}^{\infty} (u_{xx}(t, x) + |u|_x(t, x)) dx \\ &= -\xi_2'(t)\phi_* + \phi_* - u_x(t, \xi_2(t)) \\ &< \phi_* (1 - \xi_2'(t)). \end{aligned}$$

Integrating this inequality we arrive at

$$M_2(t) \leq M_2(0) + \phi_* t - (\xi_2(t) - \xi_2(0)) \phi_*.$$

On the other hand, since $u(t, \cdot) - \phi_*$ and $u(t, \cdot) - u_-(t, \cdot)$ are nonnegative by the comparison principle, it holds

$$(\xi_3(t) - \xi_2(t)) \phi_* + \int_{x_1}^{\infty} (u_-(t, x) + \phi_*) dx \leq M_2(t),$$

see also Figure 3.3. Finally, since $u_+(0, \cdot) - u(0, \cdot)$ is nonnegative, we arrive at

$$M_2(0) \leq 2x_1\phi_* + \int_{x_1}^{\infty} (u_+(0, x) + \phi_*) dx = 2x_1\phi_* + \int_0^{\infty} (\phi(x) + \phi_*) dx = \phi_*(2x_1 + 1).$$

We compute

$$\begin{aligned} F(t) &:= \int_{x_1}^{\infty} \left(\frac{u_-(t, x)}{\phi_*} + 1 \right) dx - t \\ &= \frac{1}{4} \left(-(2t + 3) \operatorname{erf} \left(\frac{x_1 - t}{2\sqrt{t}} \right) + 2x_1 \operatorname{erfc} \left(\frac{t - x_1}{2\sqrt{t}} \right) - 2e^{2t-x_1} \operatorname{erfc} \left(\frac{x_1 - 3t}{2\sqrt{t}} \right) \right. \\ &\quad \left. - 2e^{x_1} \operatorname{erfc} \left(\frac{x_1 + t}{2\sqrt{t}} \right) + e^{2(x_1+t)} \operatorname{erfc} \left(\frac{x_1 + 3t}{2\sqrt{t}} \right) + 4e^{2t-x_1} \right. \\ &\quad \left. + \frac{4\sqrt{t}}{\sqrt{\pi}} e^{-\frac{(x_1-t)^2}{4t}} - 4x_1 - 2t + 3 \right), \end{aligned}$$

and obtain

$$\lim_{t \rightarrow \infty} F(t) = \frac{3}{2} - x_1.$$

All in all, we have established

$$\phi_* (\xi_3(t) - \xi_2(t) + F(t) + t) \leq M_2(t) \leq \phi_* (2x_1 + 1 + t - (\xi_2(t) - \xi_2(0))),$$

yielding

$$\xi_3(t) - \xi_2(0) \leq 2x_1 + 1 - F(t) \rightarrow 3x_1 - \frac{1}{2} \quad \text{as } t \rightarrow \infty.$$

387 Consequently, as $x_1 < \frac{1}{6}$ there exists $t_2 > 0$ such that $\xi_2(t) < \xi_3(t) \leq \xi_2(0)$ for all $t \geq t_2$.

388 Similarly, by bounding the integral $M_1(t)$, one finds $t_1 > 0$ such that $\xi_2(0) \leq \xi_1(t) <$
 389 $\xi_2(t)$ for all $t \geq t_1$, which contradicts the fact that $\xi_2(t) < \xi_2(0)$ for all $t \geq t_2$. Hence, the
 390 interfaces $\xi_1(t)$, $\xi_2(t)$ and $\xi_3(t)$ must coalesce within finite time. \square




FIGURE 3.3. The anti-shock wave $u(t, \cdot)$ connecting the asymptotic end states $\mp\phi_*$ at $\pm\infty$, the odd subsolution $u_-(t, \cdot)$ with zero x_0 , the odd supersolution $u_+(t, \cdot)$ with zero x_1 and the interfaces $\xi_1(t)$, $\xi_2(t)$ and $\xi_3(t)$ of $u(t, \cdot)$ (we suppressed the t -dependency of the interfaces). We bound the shaded area below the graph of $u(t, \cdot)$ from below by the orange subareas.

391 4. DYNAMICS NEAR A COALESCENCE EVENT FOR SMOOTH FLUX FUNCTIONS

392 Let us consider the initial-value problem for the viscous conservation law:

$$393 \begin{cases} u_t = u_{xx} + f'(u)u_x, & t > 0, & x \in \mathbb{R}, \\ u(0, x) = u_0(x), & & x \in \mathbb{R}, \end{cases} \quad (4.1)$$

394 where $f \in C^\infty(\mathbb{R})$ satisfies $f'(0) = 0$. We assume that the initial condition $u_0 \in C^\infty(\mathbb{R})$
 395 is bounded and has bounded derivatives.

396 From the well-posedness of the viscous conservation law in the class of smooth data,
 397 cf. Lemma 2.1, we know that there exists a smooth solution $u \in C^\infty((0, \infty) \times \mathbb{R}, \mathbb{R})$ to
 398 the initial-value problem (4.1). A zero $x = \xi(t)$ of $u(t, \cdot)$ on \mathbb{R} is a C^1 -function of t as long
 399 as $u_x(t, \xi(t)) \neq 0$ by the implicit function theorem.

400 Here we classify the first two bifurcations for which the function $t \rightarrow \xi(t)$ exists for t
 401 in some interval $[0, t_0]$ with $t_0 > 0$ such that $u(t, \xi(t)) = 0$ for $t \in [0, t_0]$ and $u_x(t, \xi(t)) \neq 0$
 402 for $t \in [0, t_0)$ but may fail to exist for $t > t_0$ because we have $u_x(t_0, \xi_0) = 0$ at $\xi_0 = \xi(t_0)$.

403 **4.1. Fold bifurcation.** The main result is given by the following proposition.

Proposition 4.1. *Assume that there exists $(t_0, \xi_0) \in (0, \infty) \times \mathbb{R}$ such that*

$$u_x(t_0, \xi_0) = 0 \quad \text{and} \quad u_{xx}(t_0, \xi_0) \neq 0.$$

404 *Then, there exist two roots of $u(t, \cdot)$ near ξ_0 for $t < t_0$ near t_0 , denoted by $\xi_{1,2}(t)$, such*
 405 *that*

$$406 \quad \xi_{1,2}(t) - \xi_0 = \pm \sqrt{2(t_0 - t)} + \mathcal{O}(t_0 - t) \quad \text{as} \quad t \rightarrow t_0^- \quad (4.2)$$

407 *and*

$$408 \quad u_x(t, \xi_{1,2}(t)) = \pm \sqrt{2(t_0 - t)} u_{xx}(t_0, \xi_0) + \mathcal{O}(t_0 - t) \quad \text{as} \quad t \rightarrow t_0^-. \quad (4.3)$$

409 *No roots of $u(t, \cdot)$ near ξ_0 exist for $t > t_0$ near t_0 .*

Proof. By using the equation of motion in (4.1), we have

$$u_t(t_0, \xi_0) = u_{xx}(t_0, \xi_0) \neq 0.$$

Moreover, using Taylor expansions for smooth solutions, we obtain for any root $\xi(t)$ of $u(t, \cdot)$ near ξ_0 :

$$\begin{aligned} 0 &= u(t, \xi(t)) \\ &= \underbrace{u(t_0, \xi_0)}_{=0} + (t - t_0) \underbrace{u_t(t_0, \xi_0)}_{\neq 0} + (\xi(t) - \xi_0) \underbrace{u_x(t_0, \xi_0)}_{=0} \\ &\quad + \frac{1}{2}(t - t_0)^2 u_{tt}(t_0, \xi_0) + (t - t_0)(\xi(t) - \xi_0) u_{tx}(t_0, \xi_0) + \frac{1}{2}(\xi(t) - \xi_0)^2 \underbrace{u_{xx}(t_0, \xi_0)}_{\neq 0} + \mathcal{O}(3). \end{aligned}$$

It follows from the Newton's polygon in Figure 4.1 that this expansion defines two roots for $\xi(t)$, denoted by $\xi_{1,2}(t)$, which are given by the expansion

$$\begin{aligned} \xi_{1,2}(t) - \xi_0 &= \pm \sqrt{\frac{2u_t(t_0, \xi_0)}{u_{xx}(t_0, \xi_0)}(t_0 - t)} + \mathcal{O}(t_0 - t) \\ &= \pm \sqrt{2(t_0 - t)} + \mathcal{O}(t_0 - t), \end{aligned}$$

which exist for $t < t_0$ near t_0 , coalesce at $t = t_0$ and disappear for $t > t_0$. We also obtain

$$\begin{aligned} u_x(t, \xi_{1,2}(t)) &= \underbrace{u_x(t_0, \xi_0)}_{=0} + (t - t_0) u_{tx}(t_0, \xi_0) + (\xi_{1,2}(t) - \xi_0) \underbrace{u_{xx}(t_0, \xi_0)}_{\neq 0} + \mathcal{O}(2) \\ &= \pm \sqrt{2(t_0 - t)} u_{xx}(t_0, \xi_0) + \mathcal{O}(t_0 - t). \end{aligned}$$

410 Both expansions prove the validity of (4.2) and (4.3). □

411 **4.2. Pitchfork bifurcation.** The main result is given by the following proposition.

Proposition 4.2. *Assume that there exists $(t_0, \xi_0) \in (0, \infty) \times \mathbb{R}$ such that*

$$u_x(t_0, \xi_0) = 0, \quad u_{xx}(t_0, \xi_0) = 0, \quad \text{and} \quad u_{xxx}(t_0, \xi_0) \neq 0.$$

412 *Then, there exist three roots of $u(t, \cdot)$ near ξ_0 for $t < t_0$ near t_0 and one root near ξ_0 for*
 413 *$t > t_0$ near t_0 . Two roots, denoted by $\xi_{1,2}(t)$, are not continued for $t > t_0$ and satisfy*

$$414 \quad \xi_{1,2}(t) - \xi_0 = \pm \sqrt{6(t_0 - t)} + \mathcal{O}(t_0 - t) \quad \text{as} \quad t \rightarrow t_0^-, \quad (4.4)$$

415 *whereas the third root, denoted by $\xi(t)$, is continued for $t > t_0$ and satisfies*

$$416 \quad \xi(t) - \xi_0 = \frac{u_{tt}(t_0, \xi_0)}{2u_{xxx}(t_0, \xi_0)}(t_0 - t) + \mathcal{O}((t_0 - t)^2) \quad \text{as} \quad t \rightarrow t_0. \quad (4.5)$$

417 *We also have*

$$418 \quad u_x(t, \xi_{1,2}(t)) = 2u_{xxx}(t_0, \xi_0)(t_0 - t) + \mathcal{O}((t_0 - t)^{3/2}) \quad \text{as} \quad t \rightarrow t_0^- \quad (4.6)$$

419 *and*

$$420 \quad u_x(t, \xi(t)) = u_{xxx}(t_0, \xi_0)(t - t_0) + \mathcal{O}((t_0 - t)^2) \quad \text{as} \quad t \rightarrow t_0. \quad (4.7)$$

421 **Remark 4.3.** *The scaling laws (4.4) and (4.6) were conjectured in [13] based on numerical*
 422 *simulations of spatially odd solutions of the modular Burgers' equation. Proposition 4.2*
 423 *shows that this behavior holds for every scalar viscous conservation law (1.1) with smooth*
 424 *nonlinearity and smooth initial data.*

425 **Remark 4.4.** *The asymptotic expansions (4.4) and (4.6) imply*

$$426 \quad u_{xx}(t, \xi_{1,2}(t)) = \pm u_{xxx}(t_0, \xi_0) \sqrt{6(t_0 - t)} + \mathcal{O}(t_0 - t) \quad \text{as} \quad t \rightarrow t_0^-, \quad (4.8)$$

which was also conjectured in [13]. Indeed, if we differentiate $u(t, \xi(t)) = 0$ with the chain rule for the smooth solutions $u \in C^\infty((0, \infty) \times \mathbb{R}, \mathbb{R})$ for $t \in (0, t_0)$, while assuming that $u_x(t, \xi(t)) \neq 0$, $\xi(t)$ of $u(t, \cdot)$, then we obtain from (4.1) with $f'(0) = 0$:

$$u_t(t, \xi(t)) + \xi'(t)u_x(t, \xi(t)) = 0 \quad \Rightarrow \quad u_{xx}(t, \xi(t)) = -\xi'(t)u_x(t, \xi(t)),$$

427 *for $t \in (0, t_0)$. Hence, (4.4) and (4.6) imply (4.8). Similarly, we can derive from (4.5)*
 428 *and (4.7):*

$$429 \quad u_{xx}(t, \xi(t)) = \frac{1}{2}u_{tt}(t_0, \xi_0)(t - t_0) + \mathcal{O}((t_0 - t)^2) \quad \text{as} \quad t \rightarrow t_0, \quad (4.9)$$

430 *for the third root $\xi(t)$ which exists for all t near t_0 .*

Remark 4.5. *It follows from (4.4) and (4.5) that the three interfaces satisfy the natural ordering for the pitchfork bifurcation*

$$\xi_1(t) < \xi(t) < \xi_2(t),$$

431 *for $t < t_0$ near t_0 . It follows from (4.6) and (4.7) that the sign of the first partial derivative*
 432 *of $u(t, x)$ in x at $x = \xi(t)$ is opposite to the one at $x = \xi_{1,2}(t)$ for $t < t_0$ near t_0 .*

Remark 4.6. If $u_0(-x) = -u_0(x)$ and $f'(-z) = -f'(z)$ for $z \in \mathbb{R}$ in (4.1), then $u(t, -x) = -u(t, x)$ for every $t > 0$ and $x \in \mathbb{R}$. In this case of odd symmetry, if the assumptions of Proposition 4.2 are satisfied and $\xi_0 = 0$, then $\xi(t) = 0$ for all $t \geq 0$. Consequently, we have

$$u(t, 0) = u_{xx}(t, 0) = 0,$$

433 for all $t \geq 0$.

Proof of Proposition 4.2. By using the equation of motion in (4.1), we have

$$u_t(t_0, \xi_0) = 0 \quad \text{and} \quad u_{tx}(t_0, \xi_0) = u_{xxx}(t_0, \xi_0) \neq 0.$$

Moreover, using Taylor expansions for smooth solutions, we obtain for any root $\xi(t)$ of $u(t, \cdot)$ near ξ_0 :

$$\begin{aligned} 0 &= u(t, \xi(t)) \\ &= \underbrace{u(t_0, \xi_0)}_{=0} + (t - t_0) \underbrace{u_t(t_0, \xi_0)}_{=0} + (\xi(t) - \xi_0) \underbrace{u_x(t_0, \xi_0)}_{=0} \\ &\quad + \frac{1}{2}(t - t_0)^2 u_{tt}(t_0, \xi_0) + (t - t_0)(\xi(t) - \xi_0) \underbrace{u_{tx}(t_0, \xi_0)}_{\neq 0} + \frac{1}{2}(\xi(t) - \xi_0)^2 \underbrace{u_{xx}(t_0, \xi_0)}_{=0} \\ &\quad + \frac{1}{6}(t - t_0)^3 u_{ttt}(t_0, \xi_0) + \frac{1}{2}(t - t_0)^2 (\xi(t) - \xi_0) u_{ttx}(t_0, \xi_0) \\ &\quad + \frac{1}{2}(t - t_0)(\xi(t) - \xi_0)^2 u_{txx}(t_0, \xi_0) + \frac{1}{6}(\xi(t) - \xi_0)^3 \underbrace{u_{xxx}(t_0, \xi_0)}_{\neq 0} + \mathcal{O}(4). \end{aligned}$$

434 It follows from the Newton's polygon in Figure 4.1 that this expansion defines two sets
435 of roots. One set appears at the balance of $(t - t_0)(\xi(t) - \xi_0)$ and $(\xi(t) - \xi_0)^3$ terms and
436 the other set appears at the balance between $(t - t_0)^2$ and $(t - t_0)(\xi(t) - \xi_0)$ terms.

The former set is represented by two roots denoted as $\xi_{1,2}(t)$ which satisfy the expansion

$$\begin{aligned} \xi_{1,2}(t) - \xi_0 &= \pm \sqrt{\frac{6u_{tx}(t_0, \xi_0)}{u_{xxx}(t_0, \xi_0)}(t_0 - t) + \mathcal{O}(t_0 - t)} \\ &= \pm \sqrt{6(t_0 - t)} + \mathcal{O}(t_0 - t). \end{aligned}$$

The two roots exist for $t < t_0$ near t_0 , coalesce at $t = t_0$ and disappear for $t > t_0$. We also obtain

$$\begin{aligned} u_x(t, \xi_{1,2}(t)) &= \underbrace{u_x(t_0, \xi_0)}_{=0} + (t - t_0) \underbrace{u_{tx}(t_0, \xi_0)}_{\neq 0} + (\xi_{1,2}(t) - \xi_0) \underbrace{u_{xx}(t_0, \xi_0)}_{=0} \\ &\quad + \frac{1}{2}(t - t_0)^2 u_{ttx}(t_0, \xi_0) + (t - t_0)(\xi_{1,2}(t) - \xi_0) u_{txx}(t_0, \xi_0) \\ &\quad + \frac{1}{2}(\xi_{1,2}(t) - \xi_0)^2 \underbrace{u_{xxx}(t_0, \xi_0)}_{\neq 0} + \mathcal{O}(3), \end{aligned}$$

which implies

$$u_x(t, \xi_{1,2}(t)) = 2u_{xxx}(t_0, \xi_0)(t_0 - t) + \mathcal{O}((t_0 - t)^{3/2}) \quad \text{as } t \rightarrow t_0^-.$$

437 These expansions prove the validity of (4.4) and (4.6).

The latter set is represented by one root denoted by $\xi(t)$ which satisfies the expansion

$$\begin{aligned} \xi(t) - \xi_0 &= \frac{u_{tt}(t_0, \xi_0)}{2u_{tx}(t_0, \xi_0)}(t_0 - t) + \mathcal{O}((t_0 - t)^2) \\ &= \frac{u_{tt}(t_0, \xi_0)}{2u_{xxx}(t_0, \xi_0)}(t_0 - t) + \mathcal{O}((t_0 - t)^2). \end{aligned}$$

The root $\xi(t)$ persists for all t near t_0 . We also obtain

$$\begin{aligned} u_x(t, \xi(t)) &= \underbrace{u_x(t_0, \xi_0)}_{=0} + (t - t_0) \underbrace{u_{tx}(t_0, \xi_0)}_{\neq 0} + (\xi(t) - \xi_0) \underbrace{u_{xx}(t_0, \xi_0)}_{=0} \\ &\quad + \frac{1}{2}(t - t_0)^2 u_{ttx}(t_0, \xi_0) + (t - t_0)(\xi(t) - \xi_0) u_{txx}(t_0, \xi_0) \\ &\quad + \frac{1}{2}(\xi(t) - \xi_0)^2 \underbrace{u_{xxx}(t_0, \xi_0)}_{\neq 0} + \mathcal{O}(3), \end{aligned}$$

which implies

$$u_x(t, \xi(t)) = u_{xxx}(t_0, \xi_0)(t - t_0) + \mathcal{O}((t_0 - t)^2) \quad \text{as } t \rightarrow t_0.$$

438 These expansions prove the validity of (4.5) and (4.6). □




FIGURE 4.1. Newton's polygons used in the proofs of Proposition 4.1 (left) and Proposition 4.2 (right).

439 **4.3. Bifurcations of higher order.** By continuing the analysis from the previous two
 440 subsections, one can characterize coalescence of roots of $u(t, \cdot)$ in the non-generic case when
 441 there exists an integer $m \geq 4$ and $(t_0, \xi_0) \in (0, \infty) \times \mathbb{R}$ such that all partial derivatives
 442 of $u(t, x)$ in x at (t_0, ξ_0) are zero up to the m -th order and the m -th partial derivative of
 443 $u(t, x)$ in x at (t_0, ξ_0) is nonzero.

444 **4.4. Viscous Burgers' equation with quadratic nonlinearity.** We give a precise
 445 description of a class of solutions to the viscous Burgers' equation whose zeros undergo
 446 a pitchfork bifurcation. Thus, we take $f(u) = u^2$ in (4.1) and consider the initial value
 447 problem for the Burgers' equation

$$448 \quad \begin{cases} u_t = u_{xx} + 2uu_x, & t > 0, \\ u(0, x) = u_0(x), & x \in \mathbb{R}. \end{cases} \quad (4.10)$$

449 As is well-known, (4.10) can be solved explicitly using the Cole-Hopf transformation (see
 450 Section 3.6 in [11]). We will use the decomposition near the stationary shock-wave solution
 451 $\phi(x) = \tanh(x)$ of (4.10) to show that the pitchfork bifurcation of Proposition 4.2 does
 452 happen within finite time for all solutions of (4.10) with spatially odd initial data u_0
 453 having a single zero on $(0, \infty)$. The main result is given by the following proposition.

454 **Proposition 4.7.** *Let $u_0 \in C^\infty(\mathbb{R})$ satisfy*

- 455 • $u_0 \mp 1$, u_0' and u_0'' are L^2 -integrable on \mathbb{R}_\pm ,
- 456 • $u_0(-x) = -u_0(x)$ for $x \in \mathbb{R}$,
- 457 • for some $x_0 \in \mathbb{R}_+$, we have $u_0(x) < 0$ for $x \in (0, x_0)$ and $u_0(x) > 0$ for $x \in (x_0, \infty)$.

458 *Then, there exist a time $t_0 \in (0, \infty)$ and $\xi \in C^\infty((0, t_0), \mathbb{R}_+)$ such that the solution*
 459 *$u \in C^\infty((0, \infty) \times \mathbb{R}, \mathbb{R})$ to the initial-value problem (4.10) satisfies:*

- 460 (i) $\lim_{x \rightarrow \pm\infty} u(t, x) = \pm 1$ for $t \geq 0$,
- 461 (ii) $u(t, -x) = -u(t, x)$ for $t \geq 0$ and $x \in \mathbb{R}$,
- 462 (iii) $u(t, x) < 0$ for $x \in (0, \xi(t))$ and $u(t, x) > 0$ for $x \in (\xi(t), \infty)$ if $t \in [0, t_0)$,
- 463 (iv) $u(t, x) > 0$ for $x \in (0, \infty)$ if $t \geq t_0$.

464 *Moreover, we have $u_x(t_0, 0) = 0$, $u_{xx}(t_0, 0) = 0$, and $u_{xxx}(t_0, 0) > 0$.*

465 **Remark 4.8.** *For $u \in C^\infty((0, \infty) \times \mathbb{R}, \mathbb{R})$ and $\xi \in C^\infty((0, t_0), \mathbb{R}_+)$ in Proposition 4.7 we*
 466 *obtain the identities (4.4), (4.5), (4.6) and (4.7), since the assumptions of Proposition 4.2*
 467 *are satisfied.*

468 *Proof of Proposition 4.7.* We use the decomposition of u at the stationary shock-wave
 469 solution $x \mapsto \tanh(x)$ of $0 = 2uu_x + u_{xx}$ and write

$$470 \quad u(t, x) = \tanh(x) + v(t, x). \quad (4.11)$$

471 The perturbation v (which is not necessarily small) satisfies

$$472 \quad v_t = v_{xx} + 2vv_x + 2(\tanh(x)v)_x. \quad (4.12)$$

473 This nonlinear equation can be linearized with the Cole-Hopf transformation

$$474 \quad v(t, x) = \partial_x \log \psi(t, x). \quad (4.13)$$

475 By substituting (4.13) into (4.12), we obtain the following linear advection-diffusion equa-
476 tion

$$477 \quad \psi_t = \psi_{xx} + 2 \tanh(x) \psi_x. \quad (4.14)$$

478 We are looking for a solution of (4.14) which is bounded away from zero by a positive
479 constant. Without loss of generality, this constant can be normalized to unity, so that we
480 can look for a solution of the form

$$481 \quad \psi(t, x) = 1 + \hat{\psi}(t, x), \quad \hat{\psi}(t, \cdot) \in H^2(\mathbb{R}), \quad t \in \mathbb{R}_+. \quad (4.15)$$

482 To obtain the exact solution of (4.14), we write

$$483 \quad \hat{\psi}(t, x) = \operatorname{sech}(x) \chi(t, x) \quad (4.16)$$

484 and obtain the linear diffusion equation with constant dissipation for χ :

$$485 \quad \chi_t = \chi_{xx} - \chi. \quad (4.17)$$

486 The solutions of this linear equation are given by

$$487 \quad \chi(t, x) = \frac{e^{-t}}{\sqrt{4\pi t}} \int_{\mathbb{R}} \chi_0(y) e^{-\frac{(x-y)^2}{4t}} dy, \quad (4.18)$$

488 where $\chi_0 := \chi(0, \cdot)$ denotes the initial condition. The associated solution of the Burgers'
489 equation (4.14) is then obtained from (4.11), (4.13), (4.15), and (4.16) in the form:

$$490 \quad u(t, x) = \frac{\sinh(x) + \chi_x(t, x)}{\cosh(x) + \chi(t, x)}, \quad (4.19)$$

491 where $\chi(t, x)$ is given by (4.18).

If $\chi_0 \in C^\infty(\mathbb{R})$ satisfies $1 + \chi_0''(0) < 0$ and $\chi_0(0) > 0$, then

$$u_0'(0) = (1 + \chi_0''(0))/(1 + \chi_0(0)) < 0,$$

492 so that there exists a root $x_0 \in \mathbb{R}_+$ of u_0 . The positive root x_0 must be unique by the
493 assumptions on u_0 . Thus, we find by (4.11), (4.13), (4.15) and (4.16) that the assumptions
494 on u_0 are in one-to-one correspondence with the class of even functions $\chi_0 \in C^\infty(\mathbb{R})$ such
495 that $\operatorname{sech}(\cdot)\chi_0 \in H^2(\mathbb{R})$ and

- 496 • $\chi_0(x) > 0$ for all $x \in \mathbb{R}$,
- 497 • $x \mapsto \cosh(x) + \chi_0''(x)$ is monotonically increasing on \mathbb{R}_+ with $1 + \chi_0''(0) < 0$.

498 Now take such $\chi_0 \in C^\infty(\mathbb{R})$. Then, $\cosh(x) + \chi_0(x) > 0$ for all $x \in \mathbb{R}$ and $\sinh(x) + \chi_0'(x)$
499 has a single root $x_0 \in (0, \infty)$. Since χ_0 is even, so is $\chi \in C^\infty((0, \infty) \times \mathbb{R}, \mathbb{R})$, which
500 implies that $u(t, \cdot) \in C^\infty((0, \infty) \times \mathbb{R}, \mathbb{R})$ is spatially odd, so that (ii) holds. Furthermore,
501 $\operatorname{sech}(\cdot)\chi_0 \in H^2(\mathbb{R})$ ensures by (4.18) that $\operatorname{sech}(\cdot)\chi(t, \cdot) \in H^2(\mathbb{R})$ for all $t \geq 0$. Since
502 $\hat{\psi}(t, \cdot) \in H^2(\mathbb{R})$ for all $t \geq 0$, we have from (4.11), (4.13), and (4.15) that $\lim_{x \rightarrow \pm\infty} v(t, x) =$
503 0 and $\lim_{x \rightarrow \pm\infty} u(t, x) = \pm 1$, so that (i) holds.

It follows from the exact solution (4.18) that for every $t \geq 0$, we have $\chi(t, x) > 0$ for all $x \in \mathbb{R}$ and $x \mapsto \cosh(x) + \chi_{xx}(t, x)$ is monotonically increasing on $(0, \infty)$. Hence, $\cosh(x) + \chi(t, x) > 0$ for all $x \in \mathbb{R}$ and $\sinh(x) + \chi_x(t, x)$ has a single root $\xi(t) \in (0, \infty)$ for $t \in [0, t_0)$ as long as $1 + \chi_{xx}(t, 0) < 0$. Since

$$\chi_{xx}(t, 0) = \frac{e^{-t}}{\sqrt{4\pi t}} \int_{\mathbb{R}} \chi_0''(y) e^{-\frac{y^2}{4t}} dy$$

504 and $\operatorname{sech}(\cdot)\chi_0 \in H^2(\mathbb{R})$, the mapping $t \mapsto \chi_{xx}(t, 0)$ is monotonically increasing from a
 505 negative value $\chi_0''(0) < -1$ towards 0 as $t \rightarrow +\infty$. Hence, there exists a unique time
 506 $t_0 \in \mathbb{R}_+$ such that $1 + \chi_{xx}(t, 0)$ crosses 0 at $t = t_0$ and becomes positive for $t > t_0$ so that
 507 (iii) and (iv) hold.

Let us now show the non-degeneracy assumption at $t = t_0$ for which $u_x(t_0, 0) = 0$. Since the solution is smooth and spatially odd, we also have $u_{xx}(t_0, 0) = 0$. Since the mapping $t \mapsto \chi_{xx}(t, 0)$ is monotonically increasing and $t \mapsto \chi(t, 0)$ is monotonically decreasing, then $t \mapsto u_x(t, 0)$ is monotonically increasing, where

$$u_x(t, 0) = \frac{1 + \chi_{xx}(t, 0)}{1 + \chi(t, 0)}.$$

508 Thus, $u_{tx}(t_0, 0) > 0$ and the Burgers' equation in (4.10) implies $u_{xxx}(t_0, 0) > 0$. \square




FIGURE 4.2. An illustration of the exact solution to the Burgers' equation (4.10) obtained by means of (4.18) and (4.19). The initial condition for (4.18) is set as $\chi_0(x) := \cosh^2(1)\operatorname{sech}(x)$ so that the initial condition u_0 for (4.10) has a positive zero at $x = 1$. The integration of the exact solution in (4.18) was executed by using a numerical integration package. The root $\xi(t)$ of $u(t, \cdot)$ on $(0, \infty)$ exists for $t \in [0, t_0)$, coalesces at 0 at $t = t_0$ and disappears for $t > t_0$, where $t_0 \approx 0.205$. The solution $u(t, x)$ approaches the stationary shock wave $x \mapsto \tanh(x)$ as $t \rightarrow \infty$, which is represented by the dashed line.

509 5. NUMERICAL SIMULATIONS IN THE MODULAR BURGERS' EQUATION

510 Here we report on numerical simulations in the viscous Burgers' equation with modular
511 nonlinearity. The associated initial value problem reads

$$512 \quad \begin{cases} u_t = u_{xx} + |u|_x, & t > 0, & x \in \mathbb{R}, \\ u(0, x) = u_0(x), & & x \in \mathbb{R}. \end{cases} \quad (5.1)$$

513 Numerical computations in [13] implemented the finite-difference method for spatially
514 odd solutions of (5.1), see Lemma 3.8, for which the initial-value problem (5.1) can be
515 closed on the half-line $[0, \infty)$ subject to a Dirichlet boundary condition at $x = 0$. The
516 jump condition (3.27) was used at $x = 0$ as well as at $x = \pm\xi(t)$. The three interfaces
517 were transformed to time-independent grid points after a scaling transformation.

518 We will confirm the scaling law (1.5) of the finite-time extinction of multiple interfaces
519 in the initial-value problem (5.1). Compared to the previous numerical simulations in [13],
520 we use a regularization for the modular nonlinearity, for which the finite-difference method
521 can be implemented without any additional equations for the interface dynamics. The
522 numerical data is extracted from zeros of the solution $u(t, \cdot)$ on $(0, \infty)$ to determine the
523 power of the scaling law of the interface coalescence.

524 **5.1. Regularization.** The modular Burgers' equation can be rewritten as

$$525 \quad u_t = u_{xx} + \operatorname{sgn}(u)u_x, \quad (5.2)$$

where $\operatorname{sgn}(u)$ has a jump discontinuity at $u = 0$. To smoothen out the jump, we define
the following smooth nonlinearity for $\varepsilon > 0$,

$$f'_\varepsilon(u) := \frac{u}{\sqrt{\varepsilon^2 + u^2}}.$$

526 We have $f'_\varepsilon(u) \rightarrow \operatorname{sgn}(u)$ as $\varepsilon \rightarrow 0$ for all $u \in \mathbb{R}$, i.e. $f'_\varepsilon(u)$ converges pointwise to $\operatorname{sgn}(u)$.
527 This yields the regularized equation

$$528 \quad u_t = u_{xx} + \frac{u}{\sqrt{\varepsilon^2 + u^2}}u_x. \quad (5.3)$$

529 We consider initial data $u(0, x) = u_0(x)$ for shock and anti-shock waves with the boundary
530 condition $u_0(x) \rightarrow u_\pm$ as $x \rightarrow \pm\infty$, where u_\pm have opposite signs. The case of $u_- < 0 <$
531 u_+ includes a monotone, steadily traveling shock wave, to which the evolution of small
532 exponentially decaying perturbations converges [9]. The anti-shock case of $u_- > 0 >$
533 u_+ does not admit any steadily traveling shock-wave solutions.

534 For the simulation of shock-wave solutions with the normalized asymptotic limits
535 $u_\pm = \pm 1$, we take the following initial condition:

$$536 \quad u_0(x) = \tanh(x) \left(1 - e^{\alpha(1-x^2)} \right), \quad (5.4)$$

537 where $\alpha > 0$ is a free parameter. The parameter $\alpha > 0$ can be used to construct slopes
538 of the initial data at $x = 1$. For the simulation of anti-shock wave solutions with the

539 normalized asymptotic limits $u_{\pm} = \mp 1$, we take the negative version of (5.4), that is,

$$540 \quad u_0(x) = -\tanh(x) \left(1 - e^{\alpha(1-x^2)}\right). \quad (5.5)$$

541 Both in (5.4) and (5.5), the convergence of $u_0(x) \rightarrow u_{\pm}$ as $x \rightarrow \pm\infty$ is exponentially fast.

542 **5.2. Finite-difference method.** We rewrite the regularized Burgers' equation (5.3) in
543 the equivalent form,

$$544 \quad u_t = u_{xx} + f_{\varepsilon}(u)_x, \quad (5.6)$$

545 with $f_{\varepsilon}(u) = \sqrt{\varepsilon^2 + u_{\varepsilon}^2} - \varepsilon$.

546 We will use the Crank-Nicolson method based on the trapezoidal rule to set up our
547 numerical simulations for the equation (5.6). For the numerical discretization, we first
548 define the spatial domain $[0, L]$ partitioned into $(N + 1)$ grid points with spatial step h
549 and the time domain $[0, T]$ partitioned into M grid points with time step τ . We let x_n
550 for $0 \leq n \leq N$ be the spatial grid point and t_m for $0 \leq m \leq M$ be the temporal grid
551 point. We impose a Dirichlet condition at $x = 0$ which yields $u_0^m = 0$ and a Neumann
552 condition at $x = L$. By using the virtual grid point $x_{N+1} > L$, the Neumann condition
553 reads $u_{N+1}^m = u_{N-1}^m$.

The Crank-Nicolson method is based on the discretization rule,

$$\begin{aligned} u^{m+1} = u^m + \frac{\tau}{2h^2} [u_{n+1}^m - 2u_n^m + u_{n-1}^m + u_{n+1}^{m+1} - 2u_n^{m+1} + u_{n-1}^{m+1}] \\ + \frac{\tau}{4h} [f_{\varepsilon}(u_{n+1}^m) - f_{\varepsilon}(u_{n-1}^m) + f_{\varepsilon}(u_{n+1}^{m+1}) - f_{\varepsilon}(u_{n-1}^{m+1})]. \end{aligned}$$

We need to solve N equations for N unknowns $\{u_n^{m+1}\}_{n=1}^N$ at each $0 \leq m \leq M - 1$. Hence,
we rearrange the discretization scheme to get the unknown variables on the left and the
known variables on the right as

$$\begin{aligned} u_n^{m+1} + \frac{\tau}{h^2} u_n^{m+1} - \frac{\tau}{2h^2} (u_{n+1}^{m+1} + u_{n-1}^{m+1}) - \frac{\tau}{4h} [f_{\varepsilon}(u_{n+1}^{m+1}) - f_{\varepsilon}(u_{n-1}^{m+1})] = \\ u_n^m + \frac{\tau}{2h^2} (u_{n+1}^m + u_{n-1}^m) - \frac{\tau}{h^2} u_n^m + \frac{\tau}{4h} [f_{\varepsilon}(u_{n+1}^m) - f_{\varepsilon}(u_{n-1}^m)]. \end{aligned} \quad (5.7)$$

554 To simplify the expression, we use a predictor-corrector method (also known as Heun's
555 method). The idea is to use the solution at an initial point, u^m , and to calculate an initial
556 guess value of the next point $(u^*)^{m+1}$. Heun's method then improves this initial guess
557 value using the trapezoidal rule to determine a better estimate of the next term u^{m+1} .

To represent the predictor-corrector method, we introduce two matrices:

$$A_{\pm} = \begin{bmatrix} 1 \pm \frac{\tau}{h^2} & \mp \frac{\tau}{2h^2} & 0 & \cdots & 0 \\ \mp \frac{\tau}{2h^2} & 1 \pm \frac{\tau}{h^2} & \mp \frac{\tau}{2h^2} & \cdots & \vdots \\ \vdots & \mp \frac{\tau}{2h^2} & \ddots & \ddots & \\ \vdots & \vdots & \ddots & \ddots & \ddots \\ \vdots & \vdots & \mp \frac{\tau}{2h^2} & 1 \pm \frac{\tau}{h^2} & \mp \frac{\tau}{2h^2} \\ 0 & 0 & \cdots & \mp \frac{\tau}{h^2} & 1 \pm \frac{\tau}{h^2} \end{bmatrix},$$

where the elements of A_{\pm} at the $(N, N - 1)$ entry are doubled due to the Neumann condition $u_{N+1}^m = u_{N-1}^m$. We also represent the regularized terms in matrix vector notion,

$$b(u^m) = \begin{bmatrix} f_{\varepsilon}(u_2^m) \\ f_{\varepsilon}(u_3^m) - f_{\varepsilon}(u_1^m) \\ f_{\varepsilon}(u_4^m) - f_{\varepsilon}(u_2^m) \\ \vdots \\ f_{\varepsilon}(u_N^m) - f_{\varepsilon}(u_{N-2}^m) \\ 0 \end{bmatrix},$$

558 where we note that $f_{\varepsilon}(0) = 0$ by construction of f_{ε} and the Dirichlet condition, and
 559 $f_{\varepsilon}(u_{N+1}^m) - f_{\varepsilon}(u_{N-1}^m) = 0$ by the Neumann condition $u_{N+1}^m = u_{N-1}^m$. The correction step is
 560 computed from (5.7) by Euler's method as

$$561 \quad (u^*)^{m+1} = A_+^{-1} \left(A_- u^m + \frac{\tau}{2h} b(u^m) \right). \quad (5.8)$$

562 The prediction step is computed from (5.7) by Heun's method as

$$563 \quad u^{m+1} = A_+^{-1} \left(A_- u^m + \frac{\tau}{4h} b(u^m) + \frac{\tau}{4h} b((u^*)^{m+1}) \right). \quad (5.9)$$

We now extract the interface position $\xi(t_m)$ from u^m at $t = t_m$ by finding the two adjacent grid points x_n and x_{n+1} , where u_n and u_{n+1} are of opposite signs. By the straight line interpolation between (x_n, u_n) and (x_{n+1}, u_{n+1}) , we obtain

$$u(x) = \left(\frac{u_{n+1} - u_n}{x_{n+1} - x_n} \right) (x - x_n) + u_n.$$

564 The value of $\xi(t_m)$ is obtained by finding the root of u as

$$565 \quad \xi(t_m) = \frac{u_n x_{n+1} - u_{n+1} x_n}{u_n - u_{n+1}}. \quad (5.10)$$

566 **5.3. Numerical simulations for shock waves.** We have performed iterations on the
 567 domain $[0, L]$ discretized with the grid size $h = 0.01$. The time step was chosen to be
 568 $\tau = 0.0005$. Moreover, we took $\varepsilon = 10^{-16}$.

569 Figure 5.1 depicts the outcome of numerical simulations of the regularized approxi-
 570 mation (5.6) of the modular Burgers' equation (5.2) for the initial condition (5.4) with
 571 $\alpha = 1$ for which we take $L = 5$. It is observed that $\xi(t)$ indeed goes to 0 in finite time
 572 after which numerical computations can be continued. Yet, we stop them since we are
 573 only interested in the dynamics up to coalescence.

574 We have also performed numerical simulations for the initial condition (5.4) with $\alpha = 4$
 575 shown in Figure 5.2. For these simulations, we have taken $L = 10$ to avoid the boundary
 576 effects from the Neumann boundary condition at $x = L$. With smaller values of L , the
 577 solution decays below 1 at $x = L$ before the interface reaches 0. Although the initial
 578 condition u_0 has larger negative parts on $[0, 1]$, we observe that $\xi(t)$ still goes to 0 in a
 579 finite time. Compared to Figure 5.1, $\xi(t)$ is non-monotone as it first expands before it
 580 converges to 0.




FIGURE 5.1. Evolution of (5.6) for the initial data (5.4) with $\alpha = 1$. Left: $u(t, x)$ versus x for times $t = 0, t = 0.126$, and $t = 0.2535$. Right: evolution of $\xi(t)$ versus t .




FIGURE 5.2. The same as in Figure 5.1 but with $\alpha = 4$ and for times $t = 0, t = 0.692$, and $t = 1.3285$.

581 To confirm the scaling law (1.5) of the interface coalescence, we use linear regression
 582 in the log-log variable to approximate the associated power. That is, we consider

583
$$\log \xi(t) \text{ versus } c_1 \log(t_0 - t) + c_2, \quad (5.11)$$

584 where the coefficient c_1 represents the power of the scaling law. Note that the regres-
 585 sion (5.11) depends on the unknown time t_0 of the interface coalescence. Thus, we first
 586 conduct computations for t_0 defined on a numerical grid and obtain the best fit by mini-
 587 mizing the approximation error.




FIGURE 5.3. Left: power of the linear regression for Figure 5.1. Right: approximation error versus t_0 .




FIGURE 5.4. The same as in Figure 5.3 but for the data in Figure 5.2.

588 The outcomes of these computations are depicted in Figures 5.3 and 5.4 for the ap-
 589 proximations shown in Figures 5.1 and 5.2. The left panel shows the power versus t_0 and
 590 the right panel shows the corresponding approximation error versus t_0 . The minimal error
 591 for $\alpha = 1$ is attained at $t_0 = 0.2538$ and this value of t_0 corresponds to $c_1 = 0.5068$. The
 592 minimal error for $\alpha = 4$ is attained at $t_0 = 1.3853$ and this value of t_0 corresponds to
 593 $c_1 = 0.5127$. In both cases, the power is close to the claimed value of 0.5. We note that
 594 the time t_0 of extinction is larger for $\alpha = 4$ than for $\alpha = 1$.

595 **5.4. Numerical simulations for anti-shock waves.** We have also simulated (5.6) for
 596 the anti-shock wave initial condition (5.5). Figures 5.5 and 5.6 depict the outcomes of
 597 numerical simulations for $\alpha = 1$ and $\alpha = 4$ respectively. For $\alpha = 1$, the interface position

598 $\xi(t)$ goes to 0 monotonically, similar to the computations in Figure 5.1. For $\alpha = 4$, $\xi(t)$
 599 first expands and then reduces towards 0, similar to Figure 5.2.




FIGURE 5.5. Evolution of (5.6) for the initial data (5.5) with $\alpha = 1$. Left: $u(t, x)$ versus x for times $t = 0$, $t = 0.1635$, and $t = 0.328$. Right: evolution of $\xi(t)$ versus t .




FIGURE 5.6. The same as in Figure 5.5 but with $\alpha = 4$ and for times $t = 0$, $t = 0.7225$, and $t = 1.4455$.

600 Figures 5.7 and 5.8 show the approximate power of the scaling law and the approxi-
 601 mation error versus t_0 for the simulations shown in Figures 5.5 and 5.6. The minimum
 602 error for $\alpha = 1$ is attained at $t_0 = 0.3284$ and this value of t_0 corresponds to the power
 603 $c_1 = 0.4846$. The minimum error for $\alpha = 4$ is attained at $t_0 = 1.4459$ and this value
 604 of t_0 corresponds to $c_1 = 0.4884$. In both cases, the power is close to 0.5 and thus, the
 605 scaling law (1.5) is shown numerically to hold for anti-shock wave solutions considered
 606 here. However, the finite time of extinction is slightly larger for the anti-shock waves
 607 compared to that of the shock waves both for $\alpha = 1$ and $\alpha = 4$.




FIGURE 5.7. Left: power of the linear regression for Figure 5.5. Right: approximation error versus t_0 .




FIGURE 5.8. The same as in Figure 5.7 but for the data in Figure 5.6.

608 APPENDIX A. PROOFS OF WELL-POSEDNESS AND APPROXIMATION RESULTS

609 Here we provide proofs of the well-posedness and approximation results stated in §2.
 610 Local well-posedness of the scalar viscous conservation law (1.1), as well as approximation
 611 by solutions of the regularized equation (2.1), follows from standard theory for semilin-
 612 ear parabolic equations, cf. [10], whereas global well-posedness relies on the comparison
 613 principle, cf. [14, 17].

Proof of Lemma 2.1. First, it is well-known that ∂_x^2 is a sectorial operator on $C_{\text{ub}}(\mathbb{R})$ with domain $C_{\text{ub}}^2(\mathbb{R})$ and there exists a constant $C > 0$ such that

$$\|\partial_x^m e^{\partial_x^2 t} u\|_\infty \leq Ct^{-\frac{m}{2}} \|u\|_\infty, \tag{A.1}$$

for $m = 0, 1, 2$, $t > 0$ and $u \in C_{\text{ub}}(\mathbb{R})$. Second, the map $N: C_{\text{ub}}^1(\mathbb{R}) \rightarrow C_{\text{ub}}(\mathbb{R})$ given by $N(u) = f'(u)u_x$ is locally Lipschitz continuous since f is smooth. Third, $C_{\text{ub}}^1(\mathbb{R})$ is an intermediate space of class $J_{1/2}$ between $C_{\text{ub}}(\mathbb{R})$ and $C_{\text{ub}}^2(\mathbb{R})$. Hence, it follows from standard analytic semigroup theory, cf. [10], that there exist a maximal time $T \in (0, \infty]$ and a unique classical solution

$$u \in C([0, T], C_{\text{ub}}^1(\mathbb{R})) \cap C((0, T), C_{\text{ub}}^2(\mathbb{R})) \cap C^1((0, T), C_{\text{ub}}(\mathbb{R})),$$

614 of (1.1) with initial condition $u(0, \cdot) = u_0 \in C_{\text{ub}}^1(\mathbb{R})$. Moreover, if we have $T < \infty$, then it
 615 holds $\limsup_{t \rightarrow T^-} \|u(t, \cdot)\|_{W^{1, \infty}} = \infty$. A standard bootstrapping argument, using the fact
 616 that $f \in C^\infty(\mathbb{R})$, then yields $\partial_t^k u(t, \cdot) \in C_{\text{ub}}^l(\mathbb{R})$ for any $k, l \in \mathbb{N}_0$ and $t \in [0, T)$ implying
 617 $u \in C^\infty((0, T) \times \mathbb{R}, \mathbb{R})$.

It is well-known [14, 17] that the scalar conservation law (1.1) obeys a comparison principle yielding $m_0 \leq u(t, \cdot) \leq M_0$ for all $t \in [0, T)$ upon comparison with the constant solutions $u \equiv m_0$ and $u \equiv M_0$ of (1.1). Differentiating the mild formulation of (1.1), we obtain

$$\begin{aligned} u_x(t, \cdot) &= e^{\partial_x^2 t} u'_0 + \int_0^{t(1-\delta)} \partial_x^2 e^{\partial_x^2(t-s)} f(u(s, \cdot)) ds \\ &\quad + \int_{t(1-\delta)}^t \partial_x e^{\partial_x^2(t-s)} f'(u(s, \cdot)) u_x(s, \cdot) ds, \end{aligned} \quad (\text{A.2})$$

for $t \in [0, T)$, where $\delta \in (0, 1)$ will be fixed a posteriori. Let $R \geq 1$ be such that

$$\sup\{|f(v)| + |f'(v)| : v \in [m_0, M_0]\} \leq R.$$

Fix some $\tau \in [0, T)$. Taking norms in (A.2), while using (A.1) and the fact that $m_0 \leq u(t, \cdot) \leq M_0$, we establish

$$\begin{aligned} \|u_x(t, \cdot)\|_\infty &\leq C \|u_0\|_{W^{1, \infty}} + \int_0^{t(1-\delta)} \frac{CR}{t-s} ds + \int_{t(1-\delta)}^t \frac{CR \sup\{\|u_x(s, \cdot)\|_\infty : s \in [0, \tau]\}}{\sqrt{t-s}} ds \\ &\leq C \left(\|u_0\|_{W^{1, \infty}} + R |\log(\delta)| + 2R\sqrt{\delta t} \sup\{\|u_x(s, \cdot)\|_\infty : s \in [0, \tau]\} \right) \end{aligned}$$

for all $t \in [0, \tau]$. Thus, setting $\delta = \frac{1}{16C^2 R^2 \max\{1, \tau\}} \in (0, 1)$ and taking suprema in the latter inequality, we arrive at

$$\sup\{\|u_x(s, \cdot)\|_\infty : s \in [0, \tau]\} \leq 2C \left(\|u_0\|_{W^{1, \infty}} + R \log(16C^2 R^2 \max\{1, \tau\}) \right),$$

618 for all $\tau \in [0, T)$. We conclude that $\limsup_{t \rightarrow T^-} \|u(t, \cdot)\|_{W^{1, \infty}} = \infty$ cannot occur, implying
 619 that $T = \infty$ and the classical solution is global. \square

620 *Proof of Lemma 2.2.* Recall that ∂_x^2 is a sectorial operator on $C_{\text{ub}}(\mathbb{R})$ satisfying (A.1). In
 621 addition, the flux function $f: C_{\text{ub}}(\mathbb{R}) \rightarrow C_{\text{ub}}(\mathbb{R})$ is locally Lipschitz continuous. Hence,
 622 by a standard fixed point argument as in the proofs of [10, Theorem 7.1.2 and Proposi-
 623 tion 7.2.1] there exist a maximal time $T \in (0, \infty]$ and a unique solution $u \in C([0, T), C_{\text{ub}}(\mathbb{R}))$
 624 of (2.2). Moreover, if $T < \infty$, then it holds $\limsup_{t \rightarrow T^-} \|u(t, \cdot)\|_\infty = \infty$.

Let $\tilde{f} \in C^\infty(\mathbb{R})$ be a function satisfying

$$\sup \left\{ |f(v) - \tilde{f}(v)| : v \in [-m_0, M_0] \right\} < \delta,$$

for some $\delta > 0$. By Lemma 2.1 there exists a unique global classical solution

$$\tilde{u} \in C([0, \infty), C_{\text{ub}}^1(\mathbb{R})) \cap C((0, \infty), C_{\text{ub}}^2(\mathbb{R})) \cap C^1((0, \infty), C_{\text{ub}}(\mathbb{R}))$$

of the integral equation

$$\tilde{u}(t, \cdot) = e^{\partial_x^2 t} u_0 + \int_0^t \partial_x e^{\partial_x^2(t-s)} \tilde{f}(\tilde{u}(s, \cdot)) ds. \quad (\text{A.3})$$

satisfying $m_0 \leq \tilde{u}(t, \cdot) \leq M_0$ for all $t \geq 0$. From (2.2) and (A.3), we obtain

$$u(t, \cdot) - \tilde{u}(t, \cdot) = \int_0^t \partial_x e^{\partial_x^2(t-s)} \left(f(u(s, \cdot)) - f(\tilde{u}(s, \cdot)) + f(\tilde{u}(s, \cdot)) - \tilde{f}(\tilde{u}(s, \cdot)) \right) ds, \quad (\text{A.4})$$

for all $t \in [0, T)$. Denote by $L > 0$ the Lipschitz constant of f on $[m_0 - 1, M_0 + 1]$. Taking norms in (A.4) we arrive at

$$\|u(t, \cdot) - \tilde{u}(t, \cdot)\|_\infty \leq C \int_0^t \frac{L \|u(s, \cdot) - \tilde{u}(s, \cdot)\|_\infty + \delta}{\sqrt{t-s}} ds,$$

for any $t \in [0, T)$ with $\sup\{\|u(s, \cdot) - \tilde{u}(s, \cdot)\|_\infty : s \in [0, t]\} \leq 1$. Hence, Grönwall's Lemma [10, Lemma 7.0.3] yields a constant $M > 0$, depending only on C and L , such that

$$\|u(t, \cdot) - \tilde{u}(t, \cdot)\|_\infty \leq M\delta\sqrt{t}, \quad (\text{A.5})$$

625 for all $t \in [0, T)$ with $\sup\{\|u(s, \cdot) - \tilde{u}(s, \cdot)\|_\infty : s \in [0, t]\} \leq 1$.

We argue by contradiction and assume $T < \infty$. Take

$$0 < \delta \leq \frac{1}{2M\sqrt{T}}.$$

If

$$\sup\{\|u(s, \cdot) - \tilde{u}(s, \cdot)\|_\infty : s \in [0, T)\} > 1,$$

then by continuity, there must exist $t \in [0, T)$ with

$$\sup\{\|u(s, \cdot) - \tilde{u}(s, \cdot)\|_\infty : s \in [0, t]\} = 1.$$

However, (A.5) then implies

$$\|u(s, \cdot) - \tilde{u}(s, \cdot)\|_\infty < \frac{1}{2}$$

for any $s \in [0, t]$, which yields a contradiction. Hence, we have

$$\sup\{\|u(s, \cdot) - \tilde{u}(s, \cdot)\|_\infty : s \in [0, T)\} \leq 1$$

626 and (A.5) is satisfied for all $t \in [0, T)$. So, we must have $T = \infty$ and $u(t, \cdot)$ is global.

627 Since it holds $m_0 \leq \tilde{u}(t, \cdot) \leq M_0$ for all $t \geq 0$ and, in addition, $\delta > 0$ can be chosen
628 arbitrarily small, it follows $m_0 \leq u(t, \cdot) \leq M_0$ for all $t \geq 0$ by (A.5), which concludes the
629 proof of (2.4). \square

Proof of Lemma 2.3. First, Lemma 2.2 implies that $m_0 \leq u(t, \cdot) \leq M_0$ for all $t \geq 0$. Second, there exists by Lemma 2.2 constants $\widetilde{M}, \widetilde{\delta}_0 > 0$ such that if we take $\delta \in (0, \widetilde{\delta}_0)$, then there exists a unique global classical solution (2.3) of (2.1) satisfying $m_0 \leq \widetilde{u}(t, \cdot) \leq M_0$ and $\|u(t, \cdot) - \widetilde{u}(t, \cdot)\|_\infty \leq \widetilde{M}\delta\sqrt{t}$ for all $t \geq 0$. Thus, $\widetilde{u}(t, \cdot)$ solves the mild formulation (A.3). Subtracting (A.3) from (2.2) and differentiating we obtain

$$\begin{aligned} u_x(t, \cdot) - \widetilde{u}_x(t, \cdot) &= \int_0^{t(1-\delta)} \partial_x^2 e^{\partial_x^2(t-s)} (f(u(s, \cdot)) - f(\widetilde{u}(s, \cdot))) ds \\ &\quad + \int_0^{t(1-\delta)} \partial_x^2 e^{\partial_x^2(t-s)} \left(f(\widetilde{u}(s, \cdot)) - \widetilde{f}(\widetilde{u}(s, \cdot)) \right) ds \\ &\quad + \int_{t(1-\delta)}^t \partial_x e^{\partial_x^2(t-s)} \left(f'(u(s, \cdot)) - \widetilde{f}'(\widetilde{u}(s, \cdot)) \right) u_x(s, \cdot) ds \\ &\quad + \int_{t(1-\delta)}^t \partial_x e^{\partial_x^2(t-s)} \widetilde{f}'(\widetilde{u}(s, \cdot)) (u_x(s, \cdot) - \widetilde{u}_x(s, \cdot)) ds, \end{aligned} \tag{A.6}$$

for all $t \geq 0$. Denote by $L > 0$ the Lipschitz constant of f on $[m_0, M_0]$, and set $K = \sup\{\|u_x(s, \cdot)\|_\infty : 0 \leq s \leq \tau\}$ and $R_1 = \sup\{|f'(v)| : v \in [m_0, M_0]\}$. Thus, taking norms in (A.6), while using (A.1), we arrive at

$$\begin{aligned} \|u_x(t, \cdot) - \widetilde{u}_x(t, \cdot)\|_\infty &\leq C \int_{t(1-\delta)}^t \frac{K(R + R_1)}{\sqrt{t-s}} ds + C \int_0^t \frac{R\|u_x(s, \cdot) - \widetilde{u}_x(s, \cdot)\|_\infty}{\sqrt{t-s}} ds \\ &\quad + C \int_0^{t(1-\delta)} \frac{\delta \left(1 + L\widetilde{M}\sqrt{t}\right)}{t-s} ds, \end{aligned}$$

for all $t \in [0, \tau]$. Hence, Grönwall's Lemma [10, Lemma 7.0.3] yields a constant $M > 0$, independent of δ , such that

$$\|u_x(t, \cdot) - \widetilde{u}_x(t, \cdot)\|_\infty \leq M\sqrt{\delta t},$$

630 for all $t \geq 0$. Thus, taking $\delta_0 < \min\{\widetilde{\delta}_0, \varepsilon^2/(M^2\tau), \varepsilon/(\widetilde{M}\sqrt{\tau})\}$ we establish (2.5). \square

Proof of Lemma 2.4. We switch to the co-moving frame $\xi = x - ct$, in which equation (1.1) reads

$$w_t = w_{\xi\xi} + cw_\xi + f(w)_\xi. \tag{A.7}$$

If $w(t, \cdot)$ is a mild solution of (A.7) with initial condition $w(0, \cdot) = u_0$, then the difference $z = w - \phi$ is a mild solution of

$$z_t = (z_\xi + cz + f(z + \phi(\xi)) - f(\phi(\xi)))_\xi \tag{A.8}$$

and has initial condition $z_0 = u_0 - \phi \in C_{\text{ub}}^1(\mathbb{R}) \cap L^1(\mathbb{R})$. The integrated version of equation (A.8) reads

$$v_t = v_{\xi\xi} + cv_\xi + f(v_\xi + \phi(\xi)) - f(\phi(\xi)), \tag{A.9}$$

where the relevant solution has initial condition $v_0 \in C_{\text{ub}}^2(\mathbb{R})$ given by

$$v_0(\xi) = \int_{-\infty}^{\xi} z_0(y) dy.$$

First, the nonlinearity $N: C_{\text{ub}}^1(\mathbb{R}) \rightarrow C_{\text{ub}}(\mathbb{R})$ given by $N(v) = cv_{\xi} + f(v_{\xi} + \phi) - f(\phi)$ is well-defined and locally Lipschitz continuous. Second, ∂_{ξ}^2 is a sectorial operator on $C_{\text{ub}}(\mathbb{R})$ with dense domain $C_{\text{ub}}^2(\mathbb{R})$. Third, $C_{\text{ub}}^1(\mathbb{R})$ is an intermediate space of class $J_{1/2}$ between $C_{\text{ub}}(\mathbb{R})$ and $C_{\text{ub}}^2(\mathbb{R})$. Therefore, standard analytic semigroup theory, cf. [10, Theorem 7.1.2 and Propositions 7.1.10 and 7.2.1], yields a maximal time $T \in (0, \infty]$ and a solution $v \in C([0, T), C_{\text{ub}}^2(\mathbb{R}))$ of

$$v(t, \cdot) = e^{\partial_{\xi}^2 t} v_0 + \int_0^t e^{\partial_{\xi}^2(t-s)} (cv_{\xi}(s, \cdot) + f(v_{\xi}(s, \cdot) + \phi) - f(\phi)) ds. \quad (\text{A.10})$$

Moreover, if $T < \infty$, then we must have $\limsup_{t \rightarrow T^-} \|v(t, \cdot)\|_{W^{1, \infty}} = \infty$. Differentiating (A.10) with respect to ξ and setting $z = v_{\xi}$, we obtain

$$z(t, \cdot) = e^{\partial_{\xi}^2 t} z_0 + \int_0^t \partial_{\xi} e^{\partial_{\xi}^2(t-s)} (cz(s, \cdot) + f(z(s, \cdot) + \phi) - f(\phi)) ds. \quad (\text{A.11})$$

Hence, $z \in C([0, T), C_{\text{ub}}^1(\mathbb{R}))$ is a mild solution of (A.8) with initial condition z_0 . Thus, we have

$$v_{\xi}(t, \xi) = z(t, \xi) = w(t, \xi) - \phi = u(t, \xi + ct) - \phi(\xi),$$

where $u \in C([0, \infty), C_{\text{ub}}(\mathbb{R}))$ is the global mild solution of (A.7), established in Lemma 2.2, satisfying $\|u(t, \cdot)\|_{\infty} \leq \|u_0\|_{\infty}$ for $t \geq 0$. So, it holds

$$\|v_{\xi}(t, \cdot)\|_{\infty} = \|z(t, \cdot)\|_{\infty} \leq \|u_0\|_{\infty} + \|\phi\|_{\infty}$$

for all $t \geq 0$. Taking norms in (A.10) and using (A.1) we arrive at

$$\|v(t, \cdot)\|_{\infty} \leq C \left(\|v_0\|_{\infty} + t \sup_{0 \leq s \leq t} \|cz(s, \cdot) + f(z(s, \cdot) + \phi) - f(\phi)\|_{\infty} \right). \quad (\text{A.12})$$

631 Clearly, the right-hand side of (A.12) does not blow up as $t \rightarrow T^-$ yielding $T = \infty$. Thus,
632 we have obtained a global solution $u \in C([0, \infty), C_{\text{ub}}^1(\mathbb{R}))$ of (2.2).

Finally, we establish L^1 -integrability of $u(t, \cdot) - \phi$ for all $t \geq 0$. Since ϕ is bounded and f is locally Lipschitz continuous, we observe that the nonlinearity $G: L^1(\mathbb{R}) \cap C_{\text{ub}}(\mathbb{R}) \rightarrow L^1(\mathbb{R}) \cap C_{\text{ub}}(\mathbb{R})$ given by $G(z) = cz + f(z + \phi) - f(\phi)$ is well-defined and locally Lipschitz continuous. On the other hand, ∂_{ξ}^2 is a sectorial operator on $C_{\text{ub}}(\mathbb{R}) \cap L^1(\mathbb{R})$ and there exists a constant $C > 0$ such that

$$\|\partial_{\xi}^m e^{\partial_{\xi}^2 t} g\|_p \leq Ct^{-\frac{m}{2}} \|g\|_p, \quad (\text{A.13})$$

for $p = 1, \infty$, $m = 0, 1$, and $g \in L^p(\mathbb{R})$. Hence, by a standard fixed point argument as in the proofs of [10, Theorem 7.1.2 and Proposition 7.2.1], there exist a maximal time $\tau \in (0, \infty]$ and a unique solution $z \in C([0, \tau), C_{\text{ub}}(\mathbb{R}) \cap L^1(\mathbb{R}))$ of (A.11) such that, if $\tau < \infty$, we have $\limsup_{t \rightarrow \tau^-} \|z(t, \cdot)\|_{L^1 \cap L^{\infty}} = \infty$. Let $L > 0$ be the Lipschitz constant of

f on $[-\|u_0\| - \|\phi\|_\infty, \|u_0\|_\infty + \|\phi\|_\infty]$. Taking norms in (A.11) and using (A.13) we arrive at

$$\|z(t, \cdot)\|_1 \leq C \left(\|z_0\|_1 + \int_0^t \frac{(|c| + L)\|z(s, \cdot)\|_1}{\sqrt{t-s}} ds \right),$$

for $t \in [0, \tau)$. Hence, Grönwall's Lemma [10, Lemma 7.0.3] yields a constant $M > 0$, depending only on $C, |c|$, and L , such that

$$\|z(t, \cdot)\|_1 \leq M\|z_0\|_1,$$

633 for all $t \in [0, \tau)$. Combining the latter with $\|z(t, \cdot)\|_\infty \leq \|u_0\|_\infty + \|\phi\|_\infty$ for all $t \in [0, \tau)$
 634 yields $\tau = \infty$. We conclude that $z(t, \cdot) = w(t, \cdot) - \phi = u(t, \cdot + ct) - \phi$, and thus $u(t, \cdot) - \phi$
 635 itself, is L^1 -integrable for all $t \geq 0$. \square

636

REFERENCES

- 637 [1] S. Angenent. The zero set of a solution of a parabolic equation. *J. Reine Angew. Math.*, 390:79–96,
 638 1988.
- 639 [2] J. Crank and R. S. Gupta. A moving boundary problem arising from the diffusion of oxygen in
 640 absorbing tissue. *J. Inst. Maths. Applics*, 10:18–33, 1972.
- 641 [3] J. Endal and E. R. Jakobsen. L^1 contraction for bounded (nonintegrable) solutions of degenerate
 642 parabolic equations. *SIAM J. Math. Anal.*, 46(6):3957–3982, 2014.
- 643 [4] H. Freistühler and D. Serre. L^1 -stability of shock waves in scalar viscous conservation laws. *Comm.*
 644 *Pure Appl. Math.*, 51(3):291–301, 1998.
- 645 [5] V. A. Galaktionov and P. J. Harwin. Sturm's theorems on zero sets in nonlinear parabolic equations.
 646 In *Sturm-Liouville theory*, pages 173–199. Birkhäuser, Basel, 2005.
- 647 [6] T. Gallay and A. Scheel. Viscous shocks and long-time behavior of scalar conservation laws.
 648 *arXiv:2306.13341*, 2023.
- 649 [7] C. M. Hedberg and O. V. Rudenko. Collisions, mutual losses and annihilation of pulses in a modular
 650 nonlinear medium. *Nonlinear Dynam.*, 90(3):2083–2091, 2017.
- 651 [8] A. M. Il'in and O. A. Oleĭnik. Asymptotic behavior of solutions of the Cauchy problem for some
 652 quasi-linear equations for large values of the time. *Mat. Sb. (N.S.)*, 51(93):191–216, 1960.
- 653 [9] U. Le, D. E. Pelinovsky, and P. Pouillet. Asymptotic stability of viscous shocks in the modular
 654 Burgers equation. *Nonlinearity*, 34(9):5979–6016, 2021.
- 655 [10] A. Lunardi. *Analytic semigroups and optimal regularity in parabolic problems*, volume 16 of *Progress*
 656 *in Nonlinear Differential Equations and their Applications*. Birkhäuser Verlag, Basel, 1995.
- 657 [11] P. D. Miller. *Applied asymptotic analysis*, volume 75 of *Graduate Studies in Mathematics*. American
 658 Mathematical Society, Providence, RI, 2006.
- 659 [12] S. I. Mitchell and M. Vynnycky. The oxygen diffusion problem: Analysis and numerical solution.
 660 *Applied Math. Model.*, 39:2763–2776, 2015.
- 661 [13] D. E. Pelinovsky and B. de Rijk. Extinction of multiple shocks in the modular burgers equation.
 662 *Nonlinear Dyn.*, 111:3679–3687, 2023.
- 663 [14] M. H. Protter and H. F. Weinberger. *Maximum principles in differential equations*. Prentice-Hall,
 664 Inc., Englewood Cliffs, NJ, 1967.
- 665 [15] A. Radostin, V. Nazarov, and S. Kiyashko. Propagation of nonlinear acoustic waves in bimodular
 666 media with linear dissipation. *Wave Motion*, 50:191–195, 2013.
- 667 [16] O. V. Rudenko and C. M. Hedberg. Single shock and periodic sawtooth-shaped waves in media with
 668 non-analytic nonlinearities. *Math. Model. Nat. Phenom.*, 13(2):Paper No. 18, 27, 2018.
- 669 [17] D. Serre. L^1 -stability of nonlinear waves in scalar conservation laws. In *Evolutionary equations. Vol.*
 670 *I*, Handb. Differ. Equ., pages 473–553. North-Holland, Amsterdam, 2004.

671 [18] A. D. O. Tisbury, D. J. Needham, and A. Tzella. The evolution of travelling waves in a KPP
672 reaction–diffusion model with cut-off reaction rate. I. Permanent form travelling waves. *Stud. Appl.*
673 *Math.*, 146:301–329, 2021.

674 [19] A. D. O. Tisbury, D. J. Needham, and A. Tzella. The evolution of travelling waves in a KPP
675 reaction–diffusion model with cut-off reaction rate. II. Evolution of travelling waves. *Stud. Appl.*
676 *Math.*, 150:330–370, 2023.

677 (J. Lin) DEPARTMENT OF MATHEMATICS AND STATISTICS, MCMASTER UNIVERSITY, HAMILTON,
678 ONTARIO, CANADA, L8S 4K1

679 *Email address:* lin64@mcmaster.ca

680 (D. E. Pelinovsky) DEPARTMENT OF MATHEMATICS AND STATISTICS, MCMASTER UNIVERSITY,
681 HAMILTON, ONTARIO, CANADA, L8S 4K1 AND DEPARTMENT OF APPLIED MATHEMATICS, NIZHNY
682 NOVGOROD STATE TECHNICAL UNIVERSITY, 24 MININ STREET, 603950 NIZHNY NOVGOROD, RUSSIA

683 *Email address:* dmpeli@math.mcmaster.ca

684 (B. de Rijk) DEPARTMENT OF MATHEMATICS, KARLSRUHE INSTITUTE FOR TECHNOLOGY, EN-
685 GLERSTRASSE 2, 76131 KARLSRUHE, GERMANY

686 *Email address:* bjoern.rijk@kit.edu