

LINEAR INSTABILITY AND UNIQUENESS OF THE PEAKED PERIODIC WAVE IN THE REDUCED OSTROVSKY EQUATION*

ANNA GEYER[†] AND D. PELINOVSKY[‡]

Abstract. The stability of the peaked periodic wave in the reduced Ostrovsky equation has remained an open problem for a long time. In order to solve this problem we obtain sharp bounds on the exponential growth of the L^2 norm of co-periodic perturbations to the peaked periodic wave, from which it follows that the peaked periodic wave is linearly unstable. We also prove that the peaked periodic wave with a parabolic profile is the unique peaked wave in the space of periodic L^2 functions with zero mean and a single minimum per period.

Key words. peaked periodic wave, reduced Ostrovsky equation, characteristics, semigroup, instability

AMS subject classifications. 35B35, 35B10

DOI. 10.1137/18M117978X

1. Introduction. We address solutions of the Cauchy problem for the reduced Ostrovsky equation [33] written in the form

$$(1.1) \quad \begin{cases} u_t + uu_x = \partial_x^{-1}u, & t > 0, \\ u|_{t=0} = u_0, \end{cases}$$

where u_0 is a 2π -periodic function with zero mean defined in the Sobolev space $H_{\text{per}}^s(-\pi, \pi)$ for some $s \geq 0$, which we simply write as H_{per}^s . We denote the subspace of 2π -periodic functions with zero mean in H_{per}^s by \dot{H}_{per}^s . The operator $\partial_x^{-1} : \dot{H}_{\text{per}}^s \rightarrow \dot{H}_{\text{per}}^{s+1}$ denotes the antiderivative with zero mean, which can be defined using Fourier series.

The reduced Ostrovsky equation is also known under the names of the Ostrovsky–Hunter and Ostrovsky–Vakhnenko equation, due to contributions of Hunter [26] and Vakhnenko [43].

Local solutions to the Cauchy problem (1.1) with $u_0 \in \dot{H}_{\text{per}}^s$ exist for $s > \frac{3}{2}$ [39], and we refer to [32] for a discussion on how the well-posedness in $H^s(\mathbb{R})$ is extended to \dot{H}_{per}^s . For sufficiently large initial data, the local solutions break in finite time, similar to the inviscid Burgers equation [32]. However, if the initial data u_0 is suitably small, then the local solutions for $s = 3$ are continued for all times [19, 20]. Weak bounded solutions with shock discontinuities were constructed in [7, 8]. Weak solutions of the Cauchy problem (1.1) as the limiting solution of the Cauchy problem for the regularized Ostrovsky equation were considered in [6].

*Received by the editors April 10, 2018; accepted for publication (in revised form) January 11, 2019; published electronically April 9, 2019.

<http://www.siam.org/journals/sima/51-2/M117978.html>

Funding: The work of the second author was supported by the state task program in the sphere of scientific activity of the Ministry of Education and Science of the Russian Federation (5.5176.2017/8.9) and from the President of Russian Federation for the leading scientific schools grant NSH-2685.2018.5.

[†]Delft Institute of Applied Mathematics, Faculty Electrical Engineering, Mathematics and Computer Science, Delft University of Technology, Mekelweg 4, 2628 CD Delft, The Netherlands (A.Geyer@tudelft.nl).

[‡]Department of Mathematics and Statistics, McMaster University, Hamilton, Ontario, Canada, L8S 4K1, and Department of Applied Mathematics, Nizhny Novgorod State Technical University, 24 Minin Street, 603950 Nizhny Novgorod, Russia (dmpeli@math.mcmaster.ca).

The reduced Ostrovsky equation with smooth solutions is completely integrable as it can be reduced to the integrable Tzitzeica equation by a coordinate transformation [29]. This property enables a construction of a bi-infinite set of conserved quantities in the time evolution [5] and the inverse scattering transform with the Riemann–Hilbert approach [1]. Two integrable semidiscretizations of the reduced Ostrovsky equation have been obtained by using bilinear forms [16].

The stability of smooth and peaked periodic waves in the reduced Ostrovsky equation has been recently addressed in a number of publications [14, 18, 21, 22, 38]. By using higher-order conserved quantities the smooth small-amplitude periodic waves were shown in [14] to be unconstrained minimizers of a higher-order energy function. This result holds for *subharmonic* perturbations, that is, perturbations whose period is a multiple of the period of the smooth periodic waves. Since the higher-order conserved quantities are well-defined in the space \dot{H}_{per}^3 , where global well-posedness has been proved [20], it follows from the minimization properties that smooth small-amplitude periodic waves are both spectrally and orbitally stable. The minimization properties were confirmed numerically for smooth periodic waves of large amplitude all the way up to the limiting peaked wave of parabolic profile with maximal amplitude, for which the numerical results were inconclusive [14].

The spectral stability of smooth periodic waves with respect to *co-periodic* perturbations, that is, perturbations with the same period as the period of the periodic wave, was shown in [18] by using the standard variational formulation of the periodic waves as critical points of energy subject to fixed momentum. This result holds also for the generalized reduced Ostrovsky equation with power nonlinearity. Independently, the spectral stability of smooth periodic waves in the reduced Ostrovsky equation was shown in [22] by using a coordinate transformation of the spectral stability problem to an eigenvalue problem studied earlier in [38].

Regarding the peaked periodic waves, some conflicting results were recently obtained. In [22], the peaked wave with the parabolic profile was addressed and claimed to be “unstable in the absence of periodic boundary conditions.” A formal proof of this statement was obtained by constructing explicit solutions of the spectral stability problem for a positive (unstable) eigenvalue. However, this construction violates the periodic boundary conditions on the perturbation and hence does not provide an answer to the spectral stability question. In contrast, families of peaked periodic waves of small amplitude, which were previously unknown in the context of the reduced Ostrovsky equation, were constructed in [21] and these families were shown to be spectrally stable with respect to co-periodic perturbations by using the same coordinate transformation as in [38].

In this paper we give a simple and definite conclusion about the existence, uniqueness, and stability of peaked periodic waves in the reduced Ostrovsky equation. This is the first time, to the best of our knowledge, that the linear instability of peaked periodic waves is proved by means of semigroup theory and energy estimates.

The following theorem presents a summary of the main results of this paper. See Definitions 1, 3 and Lemmas 2, 7 for precise statements.

THEOREM 1.

1. Uniqueness: *The peaked periodic wave U_* with parabolic profile is the unique (up to spatial translations) peaked traveling wave solution of the reduced Ostrovsky equation in \dot{L}_{per}^2 having a single minimum per period. The solution is Lipschitz continuous and exists in \dot{H}_{per}^s with $s < 3/2$. Moreover, the reduced Ostrovsky equation does not admit any Hölder continuous solutions.*

2. Instability: *The orbit generated by spatial translations of the peaked periodic wave U_* is linearly unstable with respect to perturbations in X_{per}^1 , where*

$$(1.2) \quad X_{\text{per}}^1 := \{v \in \dot{L}_{\text{per}}^2 : (c_* - U_*)v \in H_{\text{per}}^1\}$$

and c_* is the wave speed of the periodic wave U_* .

Part (1) of Theorem 1 allows us to prove that the families of peaked periodic small-amplitude waves constructed in [21] do not satisfy the reduced Ostrovsky equation (see Remark 6). Our analysis relies on Fourier theory and the existence of a first integral. Indeed, the reduced Ostrovsky equation for smooth periodic waves can be rewritten as a second-order differential equation with a conserved quantity. Although this equivalence cannot be used when dealing with peaked periodic waves, we can still use a first-order invariant of the second-order differential equation to analyze the behavior of the smooth parts of the peaked periodic waves together with sharp estimates of the solution at the singularity (see Remark 4 and Lemma 2).

Part (2) of Theorem 1 gives a definite conclusion on linear instability of the peaked periodic wave with parabolic profile with respect to co-periodic perturbations. We do not make any claims regarding the spectral stability problem related to the peaked periodic wave (see Remark 19). Instead, we prove linear instability of the peaked periodic waves by obtaining sharp bounds on the exponential growth of the L^2 norm of the co-periodic perturbations in the linearized time-evolution problem in X_{per}^1 (see Lemma 7). Note that \dot{H}_{per}^1 is continuously embedded into X_{per}^1 but is not equivalent to X_{per}^1 (see Remark 8).

It is interesting to compare peaked periodic waves in the reduced Ostrovsky equation with peaked waves in other related nonlinear dispersive equations such as the Whitham equation and the Camassa–Holm equation. The existence of smooth periodic traveling waves in the Whitham equation has recently been established by [4, 10, 11, 12], where it was shown that the family of smooth periodic waves terminates at the highest, peaked wave, similarly to what happens for the reduced Ostrovsky equation. It was shown numerically in [36] that smooth periodic waves of small amplitude are stable while smooth waves of large amplitude become unstable, even before reaching the highest wave. This is different from the reduced Ostrovsky equation, where all smooth periodic waves are stable even for large amplitudes up to the peaked wave (see [14]), whereas the peaked periodic wave is unstable. For the Camassa–Holm equation, both the smooth periodic waves of all amplitudes and the limiting peaked periodic wave are stable; see [30, 31] and the earlier result [9] on peakons. It is an open question to understand which precise mechanisms govern these surprisingly different stability behaviors.

The paper is organized as follows. Section 2 contains the proof that the peaked wave with parabolic profile is unique up to spatial translations in the space of functions in \dot{L}_{per}^2 with a single minimum per period. Section 3 gives the proof of linear instability of the peaked periodic wave with respect to co-periodic perturbations. Section 4 concludes the paper with summary and discussions.

2. Peaked periodic wave. The periodic traveling waves in the reduced Ostrovsky equation are given by

$$u(x, t) = U(x - ct),$$

where c is the wave speed and U is a bounded 2π -periodic wave profile with zero

mean. The wave profile U is to be found from the boundary-value problem

$$(2.1) \quad \begin{cases} [c - U(z)]U'(z) + (\partial_z^{-1}U)(z) = 0 & \text{for every } z \in (-\pi, \pi) \text{ with } U(z) \neq c, \\ U(-\pi) = U(\pi), \quad \int_{-\pi}^{\pi} U(z)dz = 0, \end{cases}$$

where $z = x - ct$ is the traveling wave coordinate. If $U \in \dot{L}_{\text{per}}^2$, then $\partial_z^{-1}U \in \dot{H}_{\text{per}}^1$. By Sobolev's embedding, it follows that $\partial_z^{-1}U \in C_{\text{per}}$ so that the antiderivative $\partial_z^{-1}U$ with zero mean can be expressed by the pointwise formula

$$(2.2) \quad (\partial_z^{-1}U)(z) = \int_0^z U(z')dz' - \frac{1}{2\pi} \int_{-\pi}^{\pi} \int_0^z U(z')dz'dz, \quad z \in [-\pi, \pi].$$

In what follows, we assume that U is at least continuous on $[-\pi, \pi]$, that is, we assume that $U \in C_{\text{per}}$. For $\alpha \in (0, 1)$, let C_{per}^α be the space of α -Hölder 2π -periodic continuous functions such that

$$(2.3) \quad |U(x) - U(y)| \leq K|x - y|^\alpha \quad \forall x, y \in [-\pi, \pi]$$

for some $K \in \mathbb{R}$. We will adopt the following definition of single-lobe periodic waves.

DEFINITION 1. *We say that $U \in C_{\text{per}}$ is a single-lobe periodic wave if there exists $z_0 \in (-\pi, \pi)$ such that U is nonincreasing on $[-\pi, z_0]$ and nondecreasing on $[z_0, \pi]$.*

Remark 1. Due to the condition $U(-\pi) = U(\pi)$ and the symmetry of the equation

$$(c - U(z))U'(z) + \int_0^z U(z')dz' - \frac{1}{2\pi} \int_{-\pi}^{\pi} \int_0^z U(z')dz'dz = 0$$

with respect to the reflection $z \mapsto -z$, the single-lobe periodic waves in Definition 1 have even profile U with $z_0 = 0$. In this case, $(\partial_z^{-1}U)(z) = \int_0^z U(z')dz'$ is odd.

A family of smooth 2π -periodic waves to the boundary-value problem (2.1) satisfying $U(z) < c$ for every $z \in [-\pi, \pi]$ was constructed in our previous work [18] in an open interval of the speed parameter c . By Theorem 1(a) and Lemma 3 in [18], we have the following result.

LEMMA 1. *There exists $c_* > 1$ such that for every $c \in (1, c_*)$, the boundary-value problem (2.1) admits a unique smooth periodic wave in the sense of Definition 1 with the profile $U \in \dot{H}_{\text{per}}^\infty$ satisfying $U(z) < c$ for every $z \in [-\pi, \pi]$.*

Remark 2. For the smooth periodic waves $U \in \dot{H}_{\text{per}}^\infty$ to the boundary-value problem (2.1), the periodic boundary conditions are satisfied for all derivatives of U .

At $c = c_*$, the periodic wave with parabolic profile has been known since the original work of Ostrovsky [33]. It is easy to check that the boundary-value problem (2.1) is satisfied by $U(z) = (z^2 - 3c)/6$, whereas the zero mean condition is satisfied if $c = c_* := \pi^2/9$. This yields the exact expression for the peaked periodic wave with zero mean

$$(2.4) \quad U_*(z) := \frac{3z^2 - \pi^2}{18}, \quad z \in [-\pi, \pi],$$

periodically continued beyond $[-\pi, \pi]$. Note that $U_*(\pm\pi) = \pi^2/9 = c_*$ and $\pm U'_*(\pm\pi) = \frac{\pi}{3}$. The peaked periodic wave (2.4) can be represented by the Fourier cosine series

$$U_*(z) = \sum_{n=1}^{\infty} \frac{2(-1)^n}{3n^2} \cos(nz),$$




FIG. 1. The peaked wave defined in (2.4) continued periodically beyond $[-\pi, \pi]$.

which is well-defined in \dot{H}_{per}^s for $s < 3/2$. The peaked periodic wave constructed from the parabolic profile (2.4) is illustrated in Figure 1.

Remark 3. The enclosed angle at the peak of the wave is steeper than the maximal 120° angle of the Stokes wave of greatest height (see [41, 42]). Indeed, for peaked periodic waves of the reduced Ostrovsky equation the enclosed angle is $\varphi = \pi - 2\arctan(\pi/3)$, whereas for the Stokes wave it is $\varphi = \pi - 2\pi/3$.

Remark 4. The peaked periodic wave (2.4) belongs to solutions of the boundary-value problem (2.1) with profile $U_* \in \dot{H}_{\text{per}}^1$ satisfying $U_*(z) < c$ for every $z \in (-\pi, \pi)$ and $U_*(\pm\pi) = c$. The first derivative of $U_* \in \dot{H}_{\text{per}}^1$ has a finite jump singularity across the end points $z = \pm\pi$. More precisely, the profile U_* is Lipschitz continuous at $\pm\pi$, that is, there exist constants $0 < c_1 < c_2$ such that

$$c_1|z - \pi| \leq |U_*(z) - c_*| \leq c_2|z - \pi| \quad \text{for } |z - \pi| \ll 1,$$

which can be easily checked in view of the explicit expression (2.4).

The next result states that the only single-lobe periodic wave with profile $U \in C_{\text{per}}$ satisfying the boundary-value problem (2.1) and having a singularity in the derivative at $z = \pm\pi$ is the peaked periodic wave U_* given in (2.4).

LEMMA 2. *For every $c \in \mathbb{R}$, the boundary-value problem (2.1) does not admit single-lobe periodic waves in the sense of Definition 1 which are C_{per}^α with $\alpha \in [0, 1]$. The only periodic wave with a singularity in the derivative at $z = \pm\pi$ is the peaked wave with parabolic profile (2.4), which exists for $c = c_* = \pi^2/9$ and is Lipschitz at the peak.*

Proof. Let $U \in \dot{L}_{\text{per}}^2 \cap C_{\text{per}}$ be a single-lobe periodic wave solution of the boundary-value problem (2.1). By Remark 1, $U \in \dot{L}_{\text{per}}^2$ is even, $\partial_z^{-1}U \in \dot{H}_{\text{per}}^1$ is odd, and $\partial_z^{-1}U$ is represented by the Fourier sine series which converges absolutely and uniformly, so that $(\partial_z^{-1}U)(\pm\pi) = 0$.

Let us first consider the case where $U(z) \neq c$ for every $z \in (-\pi, \pi)$ and $U(\pm\pi) = c$. Let $\alpha \in (0, 1)$. We assume to the contrary that there exists a solution U of the boundary-value problem (2.1) with $U \in C_{\text{per}}^\alpha$. If $U \in C_{\text{per}}^\alpha$ with $\alpha \in (0, 1)$, then $\partial_z^{-1}U \in C_{\text{per}}^1$. Since $U(\pm\pi) = c$ and $(\partial_z^{-1}U)(\pm\pi) = 0$ we find that $c - U(z) \sim (\pi - z)^\alpha$ and $(\partial_z^{-1}U)(z) \sim (\pi - z)$ at $z = \pm\pi$. Since U satisfies the boundary value problem (2.1) we have that

$$(2.5) \quad U'(z) = -\frac{(\partial_z^{-1}U)(z)}{c - U(z)}, \quad z \in (-\pi, \pi),$$

which yields $U'(z) \sim (\pi - z)^{1-\alpha}$ at $z = \pm\pi$. Equation (2.5) also implies that $U' \in C^1(-\pi, \pi)$ so we find that $U' \in C_{\text{per}}^{1-\alpha}$. Since $1 - \alpha \in (0, 1)$ we conclude that $U \in C_{\text{per}}^1$ in contradiction to the assumption that $U \in C_{\text{per}}^\alpha$ with $\alpha \in (0, 1)$. The case $\alpha = 0$,




FIG. 2. Phase plane portrait obtained from level curves of the first-order invariant (2.6) for some $c > 0$. The dashed black line indicates the singularity line $U = c$. The solid black curve to the left of the singular line corresponds to the parabolic profile (2.4).

which refers to solutions $U \in C_{\text{per}}$ in view of Definition 1, can be proved in exactly the same way.

We now show that the only peaked periodic solution with peak at $U(\pm\pi) = c$ is the solution with the parabolic profile (2.4). Since $U(z) \neq c$ for every $z \in (-\pi, \pi)$ and $U \in C^1(-\pi, \pi)$, the first-order invariant

$$\begin{aligned}
 E &= \frac{1}{2} [c - U(z)]^2 [U'(z)]^2 + \frac{c}{2} U(z)^2 - \frac{1}{3} U(z)^3 \\
 (2.6) \quad &= \frac{1}{2} [(\partial_z^{-1}U)(z)]^2 + \frac{c}{2} U(z)^2 - \frac{1}{3} U(z)^3
 \end{aligned}$$

holds for $z \in (-\pi, \pi)$. Since $(\partial_z^{-1}U)(z)$ is continuous in $z = \pm\pi$ with $(\partial_z^{-1}U)(\pm\pi) = 0$, E is continuous and constant up to the boundary at $z = \pm\pi$ and we have $E|_{z=\pm\pi} = c^3/6 =: E_c$. For $c > 0$, the level with $E = E_c$ (see the bold curve in Figure 2) gives rise to a peaked periodic wave solution with parabolic profile $U(z) < c$. We claim that this peaked wave is exactly the solution (2.4) with speed $c = c^*$. Indeed, from the level set with $E = E_c$ we have that

$$\frac{1}{2}(c - U)^2(U'(z))^2 = \frac{c^3}{6} - \frac{c}{2}U^2 + \frac{1}{3}U^3 = \frac{1}{6}(c - U)^2(c + 2U)$$

and hence

$$U'(z) = \frac{1}{\sqrt{3}}\sqrt{c + 2U} > 0 \quad \text{for } z \in (0, \pi)$$

subject to the boundary condition $U'(0) = 0$. By separation of variables we can solve this equation uniquely to find that $U(z) = \frac{1}{6}(z^2 - 3c)$. In view of the condition $U(\pi) = c$, this implies that $c = \frac{\pi^2}{9} = c^*$, which proves the claim.

Let us now analyze the situation where there exists $z_1 \in (0, \pi)$ such that $U(\pm z_1) = c$ and (2.5) holds separately for $z \in (0, z_1)$ and for $z \in (z_1, \pi)$. Let us assume that $U(z) < c$ for $z \in (0, z_1)$. If $U(z) > c$ for $z \in (0, z_1)$, the proof is analogous. There are two possibilities, either $(\partial_z^{-1}U)(\pm z_1) = 0$ or $(\partial_z^{-1}U)(\pm z_1) \neq 0$.

If $(\partial_z^{-1}U)(\pm z_1) = 0$, then by the same argument as above the first-order invariant E is continuous and constant on $[-z_1, z_1]$ with $E|_{z=\pm z_1} = E_{z_1}$.

For $z \in (z_1, \pi]$ the solution corresponding to the level set $E = E_{z_1}$ can either be continued uniquely from the region with $U(z) < c$ into the region with $U(z) > c$, or $z = z_1$ represents a turning point (a local maximum for U) and the solution can be continued uniquely into the region with $U(z) < c$ for $z \gtrsim z_1$.¹ However, if z_1 is not a turning point, then $U(z) > 0$ for $c > 0$ and $U(z) < 0$ for $c < 0$ in the interval $z \in (z_1, \pi]$. Therefore, the first variant implies that $(\partial_z^{-1}U)(z) = \int_0^z U(z')dz' \neq 0$ for every $z \in (z_1, \pi]$, which contradicts $(\partial_z^{-1}U)(\pi) = 0$. If z_1 is a turning point, then U does not belong to the class of single-lobe periodic waves (see Remark 5).

If $(\partial_z^{-1}U)(\pm z_1) \neq 0$, then the contradiction arises from the fact that, since $(\partial_z^{-1}U)(z)$ is continuous at z_1 and (2.5) holds separately for $z \in (0, z_1)$ and for $z \in (z_1, \pi)$, the change of the sign of $U'(z)$ across z_1 is determined by the change of the sign of $c - U(z)$ across z_1 . Indeed, if $U(z) < c$ both for $z \lesssim z_1$ and $z \gtrsim z_1$, then it follows from (2.5) that the sign of $U'(z)$ remains the same for $z \in (0, z_1)$ and $z \in (z_1, \pi)$. But this is impossible since $U'(z)$ must change sign for $z \lesssim z_1$ and $z \gtrsim z_1$ if $U(z) < c$ on both sides of z_1 . If on the other hand $U(z) < c$ for $z \lesssim z_1$ and $U(z) > c$ for $z \gtrsim z_1$, then it follows again from (2.5) that the sign of $U'(z)$ changes, in contradiction with the monotone increase of $U(z)$ for all $z \in (0, \pi)$. Hence, both possibilities with $(\partial_z^{-1}U)(\pm z_1) \neq 0$ yield a contradiction.

Combining all these arguments we find that the only single-lobe peaked periodic wave has a parabolic profile (2.4), which is Lipschitz at the peak $U(\pm\pi) = c$. \square

Remark 5. There is a simple way to obtain other peaked periodic waves in the boundary-value problem (2.1). One can flip the periodic wave with a parabolic profile at a point $z_0 \in (0, \pi)$ and pack two such waves over one period. This possibility is allowed in the proof of Lemma 2, but not in the class of single-lobe periodic waves. Similarly, one can pack three or more periods of the peaked wave with parabolic profiles. Definition 1 eliminates this type of nonuniqueness of the peaked periodic wave in the boundary-value problem (2.1).

Remark 6. With the following formal transformation,

$$(2.7) \quad U(z) = \mathbf{u}(\zeta), \quad z = \int_0^\zeta (c - \mathbf{u}(\zeta'))d\zeta',$$

smooth periodic waves with profile U satisfying the quasilinear second-order equation²

$$(2.8) \quad \frac{d}{dz} [c - U(z)] \frac{dU}{dz} + U = 0$$

are related to smooth periodic waves with profile \mathbf{u} satisfying the semilinear second-order equation

$$(2.9) \quad \frac{d^2 \mathbf{u}}{d\zeta^2} + (c - \mathbf{u})\mathbf{u} = 0.$$

Although all periodic solutions of (2.9) are smooth, the coordinate transformation (2.7) fails to be invertible if $\mathbf{u}(\zeta) = c$ for some ζ . Such points generate singularities in the periodic solutions of the quasi-linear equation (2.8) since

$$U'(z) = \frac{\mathbf{u}'(\zeta)}{c - \mathbf{u}(\zeta)}.$$

¹The notation $z \gtrsim z_1$ means that $0 < z - z_1 < \varepsilon$ for some small $\varepsilon > 0$, and equivalently for the reverse inequality.

²The quasi-linear equation (2.8) is a derivative of the first equation in the boundary-value problem (2.1) in z , which is justified for smooth periodic waves in Lemma 1.

In [21], small-amplitude peaked periodic waves of (2.8) with $c < 0$ were constructed from small-amplitude smooth periodic waves of (2.9) with a coordinate transformation similar to (2.7). The corresponding profile U for such peaked periodic waves has a square root singularity of the form

$$(2.10) \quad U(z) = c + \mathcal{O}(\sqrt{\pi^2 - z^2}) \quad \text{as } z \rightarrow \pm\pi.$$

Our analysis in the proof of Lemma 2 shows³ that such solutions cannot exist, since the expansion (2.10) implies that $U \in C_{\text{per}}^{1/2}$. We conclude that the small-amplitude peaked periodic waves constructed in [21] do not satisfy the boundary-value problem (2.1). A similar conclusion follows from the recent work in [4].

Remark 7. In [40] several different types of traveling wave solutions to the reduced Ostrovsky equation were constructed by means of phase plane analysis. Three types of solitary waves (see Figure 9 in [40]) were found for $c < 0$. One of them is a loop soliton, given by a multivalued function, which is studied in many publications [15, 43, 44]. The other two solutions have points of infinite slope (cusps), either at the maximum or at the inflection points. The cusped solitary waves were also constructed in [38] by using the transformation (2.7). By using similar arguments as in the proof of Lemma 2, the cusped waves are proved not to be weak solutions to the reduced Ostrovsky equation.

3. Linear instability of the peaked periodic wave. We add a co-periodic perturbation v to the traveling wave U , that is, a perturbation with the same period 2π . Truncating the quadratic terms and moving with the reference frame of the traveling wave yields the linearized evolution problem in the form

$$(3.1) \quad \begin{cases} v_t + \partial_z [(U(z) - c)v] = \partial_z^{-1} v, & t > 0, \\ v|_{t=0} = v_0. \end{cases}$$

The linearized evolution equation can be formulated in the form $v_t = \partial_z Lv$ defined by the self-adjoint operator

$$(3.2) \quad L = P_0 (\partial_z^{-2} + c - U(z)) P_0 : \dot{L}_{\text{per}}^2 \rightarrow \dot{L}_{\text{per}}^2,$$

where $P_0 : L_{\text{per}}^2 \rightarrow \dot{L}_{\text{per}}^2$ is the projection operator that removes the mean value of 2π -periodic functions. The form $v_t = \partial_z Lv$ is related to the formulation of the reduced Ostrovsky equation in the traveling wave coordinate $z = x - ct$ as a Hamiltonian system defined by the symplectic operator ∂_z and the conserved energy function $H_c(u) = H(u) + cQ(u)$, where

$$(3.3) \quad H(u) = \int_{-\pi}^{\pi} \left[-(\partial_z^{-1} u)^2 - \frac{1}{3} u^3 \right] dz, \quad Q(u) = \int_{-\pi}^{\pi} u^2 dz$$

are the conserved energy and momentum functionals for the reduced Ostrovsky equation (1.1). The periodic wave $u = U$ is a critical point of $H_c(u)$ and the self-adjoint operator L is the Hessian operator of the energy function $H_c(u)$ at the periodic wave $u = U$.

³Solutions of [21] have nonzero mean value, and hence Lemma 2 does not apply directly. However, the arguments in the proof lead to the same conclusion also for solutions with nonzero mean. Indeed, if U has a nonzero mean, $\partial_z^{-1} U(z)$ may not be zero at $z = \pm\pi$. However, if we translate the solution by half a period so that the singularity is placed at $z = 0$, then $\partial_z^{-1} U(0) = 0$ by oddness of $\partial_z^{-1} U$ and we can use the same contradiction as the one obtained from (2.5).

Thanks to the translational invariance of the boundary-value problem (2.1), $L\partial_z U = 0$, where $\partial_z U \in \dot{L}^2_{\text{per}}$, holds for both the smooth periodic waves of Lemma 1 and the peaked periodic wave (2.4) in Lemma 2. Associated to the translational eigenvector is the symplectic orthogonality constraint $\langle U, v \rangle = 0$. This constraint (see [3, 24, 34]) is used to study both the evolution of the Cauchy problem (3.1) and the spectrum of the linearized operator

$$(3.4) \quad \partial_z L : X^1_{\text{per}} \subset \dot{L}^2_{\text{per}} \rightarrow \dot{L}^2_{\text{per}},$$

where $X^1_{\text{per}} = \{v \in \dot{L}^2_{\text{per}} : (c - U)v \in H^1_{\text{per}}\}$ is the maximal domain of $\partial_z L$.

Remark 8. For smooth periodic waves we have $c - U(z) > 0$ for every $z \in [-\pi, \pi]$ so that $X^1_{\text{per}} \equiv \dot{H}^1_{\text{per}}$. For the peaked periodic wave U_* with speed c_* , the space \dot{H}^1_{per} is continuously embedded into X^1_{per} since U_* is bounded, but \dot{H}^1_{per} is not equivalent to X^1_{per} . Indeed, if a perturbation v to U_* is piecewise C^1_{per} with a finite jump-discontinuity at $z = \pm\pi$, then $v \notin \dot{H}^1_{\text{per}}$ but $v \in X^1_{\text{per}}$ since $(c_* - U_*)v \in \dot{H}^1_{\text{per}}$ in view of the fact that $U_*(\pm\pi) = c_*$.

In what follows, $\langle \cdot, \cdot \rangle$ and $\|v\|_{L^2_{\text{per}}}$ denote the inner product and the L^2 norm with integration over $[-\pi, \pi]$, respectively. In the case of the peaked periodic wave U_* with the speed c_* , we equip X^1_{per} with the norm

$$(3.5) \quad \|v\|_{X^1_{\text{per}}} := \|v\|_{L^2_{\text{per}}} + \|\partial_z [(c_* - U_*)v]\|_{L^2_{\text{per}}}.$$

We distinguish two concepts of stability of 2π -periodic waves with respect to linearization.

DEFINITION 2. *The traveling wave U is said to be spectrally stable if $\sigma(\partial_z L) \subset i\mathbb{R}$ in \dot{L}^2_{per} . Otherwise, it is said to be spectrally unstable.*

DEFINITION 3. *The traveling wave U is said to be linearly stable if for every $v_0 \in X^1_{\text{per}}$ satisfying $\langle U, v_0 \rangle = 0$, there exists $C > 0$ and a unique global solution $v \in C(\mathbb{R}, X^1_{\text{per}})$ to the Cauchy problem (3.1) such that*

$$(3.6) \quad \|v(t)\|_{X^1_{\text{per}}} \leq C \|v_0\|_{X^1_{\text{per}}}, \quad t > 0.$$

Otherwise, it is said to be linearly unstable.

In [18], we have proved that the smooth periodic waves of Lemma 1 are spectrally stable in the sense of Definition 2. Here we intend to show that the peaked periodic wave U_* of Lemma 2 given in (2.4) is linearly unstable in the sense of Definition 3. The linear instability is due to the sharp exponential growth of the unique global solution to the Cauchy problem (3.1) with $U = U_*$:

$$(3.7) \quad C \|v_0\|_{L^2_{\text{per}}} e^{\pi t/6} \leq \|v(t)\|_{L^2_{\text{per}}} \leq \|v_0\|_{L^2_{\text{per}}} e^{\pi t/6}, \quad t > 0,$$

for some $C \in (0, 1)$. We will obtain these bounds in two steps. In the first step, carried out in section 3.1, we apply the method of characteristics to the truncated linearized equation (3.1) without the dispersive term $\partial_z^{-1}v$ and obtain the sharp bounds (3.7) for all initial conditions $v_0 \in X^1_{\text{per}}$ satisfying the constraint

$$(3.8) \quad \int_{-\pi}^{\pi} z v_0(z)^2 dz = 0.$$

In the second step, carried out in section 3.2, we will show that the bounds (3.7) remain true in the full linearized equation (3.1) for a subset of initial conditions $v_0 \in X_{\text{per}}^1$ satisfying the constraint (3.8) and the additional constraint

$$(3.9) \quad \int_{-\pi}^{\pi} z^2 v_0(z) dz = 0,$$

which arises due to the orthogonality condition $\langle U, v \rangle = 0$ in Definition 3 and the zero-mean condition on v_0 . Regarding spectral stability or instability of the peaked periodic wave (2.4), we will show in section 3.3 that $\sigma(L)$ in \dot{L}_{per}^2 is given by a continuous spectrum on $[0, \pi^2/6]$, which includes the embedded eigenvalue $\lambda_0 = 0$ with the eigenvector $\partial_z U$, and a simple negative eigenvalue $\lambda_1 < 0$. As a result, no spectral gap appears between $\lambda_0 = 0$ and the continuous spectrum, and hence it is impossible to solve the spectral stability problem by applying the standard methods from [3, 24, 34].

3.1. Linear instability of truncated evolution. For the peaked periodic wave (2.4), we obtain the simple expression

$$(3.10) \quad U_*(z) - c_* = \frac{1}{6}(z^2 - \pi^2), \quad z \in [-\pi, \pi].$$

By removing the term $\partial_z^{-1} v$ from the linearized evolution problem (3.1) and using the explicit expression (3.10), we can write the truncated evolution problem in the form

$$(3.11) \quad \begin{cases} v_t + \frac{1}{6} \partial_z [(z^2 - \pi^2)v] = 0, & t > 0, \\ v|_{t=0} = v_0, \end{cases}$$

where the initial data v_0 is taken in X_{per}^1 . The evolution problem can be solved by the method of characteristics along the family of characteristic curves $z = Z(s, t)$, where $s \in [-\pi, \pi]$ is a parameter for the initial data and $t \geq 0$ is the evolution time. Defining

$$(3.12) \quad \begin{cases} \frac{d}{dt} Z(s, t) = \frac{1}{6} [Z(s, t)^2 - \pi^2], & t > 0, \\ Z(s, 0) = s \end{cases}$$

and setting $V(s, t) := v(Z(s, t), t)$ yields the evolution problem in the form

$$(3.13) \quad \begin{cases} \frac{d}{dt} V(s, t) = -\frac{1}{3} Z(s, t) V(s, t), & t > 0, \\ V(s, 0) = v_0(s). \end{cases}$$

The family of characteristic curves is obtained by integrating the differential equation (3.12) with the parameter $s \in [-\pi, \pi]$. Because $Z = \pm\pi$ are critical points of the differential equation (3.12), the family of characteristic curves remain inside the invariant region $[-\pi, \pi]$ for every $t \geq 0$. The family of characteristic curves can be obtained in the explicit form

$$(3.14) \quad Z(s, t) = \pi \frac{s \cosh(\pi t/6) - \pi \sinh(\pi t/6)}{\pi \cosh(\pi t/6) - s \sinh(\pi t/6)}, \quad s \in [-\pi, \pi], \quad t \in \mathbb{R}.$$

For later use of the chain rule we compute

$$(3.15) \quad \frac{\partial}{\partial s} Z(s, t) = \frac{\pi^2}{[\pi \cosh(\pi t/6) - s \sinh(\pi t/6)]^2}, \quad s \in [-\pi, \pi], \quad t \in \mathbb{R}.$$

The explicit solution for V in characteristic variables is obtained by integrating the differential equation (3.13) with respect to the parameter $s \in [-\pi, \pi]$:

$$V(s, t) = v_0(s)e^{-\frac{1}{3} \int_0^t Z(s, t') dt'}.$$

In view of (3.15), the explicit solution is given by

$$(3.16) \quad V(s, t) = \frac{1}{\pi^2} [\pi \cosh(\pi t/6) - s \sinh(\pi t/6)]^2 v_0(s), \quad s \in [-\pi, \pi], \quad t \in \mathbb{R}.$$

Remark 9. Since $Z(\pm\pi, t) = \pm\pi$ for every $t \in \mathbb{R}$, we have $V(\pm\pi, t) = e^{\mp\pi t/3} v(\pm\pi, t)$, and hence $V(-\pi, t) = V(\pi, t)$ for $t \neq 0$ if and only if $v_0(\pm\pi) = 0$, in which case $V(\pm\pi, t) = 0$ for every $t \in \mathbb{R}$. Therefore, $V(\cdot, t) \notin \dot{H}_{\text{per}}^1$ for $t \neq 0$ if $v_0 \in \dot{H}_{\text{per}}^1$ with $v_0(\pm\pi) \neq 0$.

By using the explicit solutions (3.14) and (3.16), we are able to state and prove the following linear instability result for the truncated evolution problem (3.11).

LEMMA 3. *For every $v_0 \in X_{\text{per}}^1$, there exists a unique global solution $v \in C(\mathbb{R}, X_{\text{per}}^1)$ to the Cauchy problem (3.11) satisfying the upper bound*

$$(3.17) \quad \|v(t)\|_{L_{\text{per}}^2} \leq \|v_0\|_{L_{\text{per}}^2} e^{\pi t/6}, \quad t > 0.$$

If $\int_{-\pi}^{\pi} s v_0(s)^2 ds = 0$, then the global solution satisfies the lower bound

$$(3.18) \quad \frac{1}{2} \|v_0\|_{L_{\text{per}}^2} e^{\pi t/6} \leq \|v(t)\|_{L_{\text{per}}^2}, \quad t > 0.$$

Proof. The existence of a global solution in the explicit form (3.14) and (3.16) is obtained from the method of characteristics. By using the chain rule and (3.15), we verify that the mean-zero constraint is preserved by the time evolution:

$$\int_{-\pi}^{\pi} v(z, t) dz = \int_{-\pi}^{\pi} V(s, t) \frac{\partial Z}{\partial s} ds = \int_{-\pi}^{\pi} v_0(s) ds = 0.$$

The explicit expression (3.16) implies that $V(\cdot, t) \in X_{\text{per}}^1$ if $v_0 \in X_{\text{per}}^1$ and $t \in \mathbb{R}$. On the other hand, the explicit expression (3.14) implies that for every $\tau > 0$, there exists $C_\tau > 0$ such that

$$\frac{\partial}{\partial s} Z(s, t) \geq C_\tau, \quad s \in [-\pi, \pi], \quad t \in [-\tau, \tau].$$

Hence, the chain rule implies that $v(\cdot, t) \in X_{\text{per}}^1$ if $v_0 \in X_{\text{per}}^1$ and $t \in \mathbb{R}$. The uniqueness of such global solutions follows by standard theory (see Theorem 3.1 in [2]).

It remains to prove the sharp exponential growth in the bounds (3.17) and (3.18). By the chain rule, we obtain

$$\begin{aligned} \int_{-\pi}^{\pi} v(z, t)^2 dz &= \int_{-\pi}^{\pi} V(s, t)^2 \frac{\partial Z}{\partial s} ds \\ &= \frac{1}{\pi^2} \int_{-\pi}^{\pi} [\pi \cosh(\pi t/6) - s \sinh(\pi t/6)]^2 v_0(s)^2 ds. \end{aligned}$$

From here, we have the upper bound

$$\|v(t)\|_{L_{\text{per}}^2}^2 \leq e^{\pi t/3} \|v_0\|_{L_{\text{per}}^2}^2$$

and the lower bound under the additional condition $\int_{-\pi}^{\pi} s v_0(s)^2 ds = 0$:

$$\|v(t)\|_{L^2_{\text{per}}}^2 = \cosh(\pi t/6)^2 \|v_0\|_{L^2_{\text{per}}}^2 + \frac{1}{\pi^2} \sinh(\pi t/6)^2 \|s v_0\|_{L^2_{\text{per}}}^2 \geq \frac{1}{4} e^{\pi t/3} \|v_0\|_{L^2_{\text{per}}}^2.$$

Taking the square root of these bounds yields (3.17) and (3.18). □

Remark 10. By the chain rule, we also have

$$\begin{aligned} & \int_{-\pi}^{\pi} [\partial_z(\pi^2 - z^2)v(z, t)]^2 dz \\ &= \frac{1}{\pi^2} \int_{-\pi}^{\pi} [\pi \cosh(\pi t/6) - s \sinh(\pi t/6)]^2 [\partial_s(\pi^2 - s^2)v_0(s)]^2 ds, \end{aligned}$$

from which the sharp exponential growth with the same growth rate as in the bounds (3.17) and (3.18) can be established for the second term in the X^1_{per} norm given by (3.5).

Remark 11. The global solution in Lemma 3 remains bounded in L^1 . This follows from the chain rule:

$$\int_{-\pi}^{\pi} |v(z, t)| dz = \int_{-\pi}^{\pi} |V(s, t)| \frac{\partial Z}{\partial s} ds = \int_{-\pi}^{\pi} |v_0(s)| ds.$$

Since

$$(3.19) \quad \|v_0\|_{L^1_{\text{per}}} \leq (2\pi)^{1/2} \|v_0\|_{L^2_{\text{per}}},$$

$v_0 \in \dot{L}^2_{\text{per}}$ implies $v_0 \in L^1_{\text{per}}$. Extending this bound to the time-dependent solution,

$$(3.20) \quad \|v(t)\|_{L^1_{\text{per}}} \leq (2\pi)^{1/2} \|v(t)\|_{L^2_{\text{per}}}, \quad t > 0,$$

shows that the L^1 norm of the global solution $v(t)$ may remain bounded even if the L^2 norm of this solution grows exponentially.

Remark 12. Truncating a quadratic form associated with the self-adjoint operator L in (3.2) and using the chain rule yields the energy conservation for the truncated evolution (3.11):

$$\int_{-\pi}^{\pi} (\pi^2 - z^2)v(z, t)^2 dz = \int_{-\pi}^{\pi} [\pi^2 - Z(s, t)^2] V(s, t)^2 \frac{\partial Z}{\partial s} ds = \int_{-\pi}^{\pi} (\pi^2 - s^2)v_0(s)^2 ds.$$

The energy conservation shows that the truncated evolution leads to the exponential growth of $\|v(t)\|_{L^2_{\text{per}}}^2$ and $\|zv(t)\|_{L^2_{\text{per}}}^2$ but the difference between the two squared norms remains bounded.

Remark 13. For the smooth periodic waves of Lemma 1 satisfying $U(z) < c$ for every $z \in [-\pi, \pi]$, the truncated energy $\int_{-\pi}^{\pi} (c - U)v^2 dz$ is coercive in the L^2 norm, and hence the energy conservation

$$\int_{-\pi}^{\pi} [c - U(z)] v(z, t)^2 dz = \int_{-\pi}^{\pi} [c - U(z)] v_0(s)^2 ds$$

implies a global time-independent bound on $\|v(t)\|_{L^2_{\text{per}}}^2$, where $v(t)$ is a solution of the truncation of the linear evolution equation (3.1) without the $\partial_z^{-1}v$ term.

Remark 14. For the smooth periodic waves of Lemma 1, the characteristic curves reach the boundaries $z = \pm\pi$ in finite time because $z = \pm\pi$ are not critical points of the differential equations for the characteristic curves. On the other hand, for the peaked periodic wave (2.4), the characteristic curves reach the boundaries $z = \pm\pi$ in infinite time. The latter property induces exponential growth of the global solutions to the Cauchy problem (3.11) as is shown in Lemma 3.

3.2. Linear instability of full evolution. Here we consider the full linearized evolution problem (3.1) with (3.10) and rewrite the evolution problem in the form

$$(3.21) \quad \begin{cases} v_t + \frac{1}{6}\partial_z [(z^2 - \pi^2)v] = \partial_z^{-1}v, & t > 0, \\ v|_{t=0} = v_0, \end{cases}$$

where the initial data v_0 is taken in X_{per}^1 .

LEMMA 4. *For every $v_0 \in X_{\text{per}}^1$ there exists a unique global solution $v \in C(\mathbb{R}, X_{\text{per}}^1)$ of the Cauchy problem (3.21).*

Proof. By Lemma 3, the Cauchy problem (3.11) with $v_0 \in X_{\text{per}}^1$ has a unique global solution $v \in C(\mathbb{R}, X_{\text{per}}^1)$. In the framework of semigroup theory, the evolution equation (3.11) can be written in the form $v_t = A_0v$, where

$$A_0 := \frac{1}{6}\partial_z(\pi^2 - z^2)v.$$

The existence of a unique global solution $v \in C(\mathbb{R}, X_{\text{per}}^1)$ implies that the operator A_0 with domain $D(A_0) = X_{\text{per}}^1$ is the infinitesimal generator of a strongly continuous semigroup $(S_0(t))_{t \geq 0}$ on \dot{L}_{per}^2 . Since $\partial_z^{-1} : \dot{L}_{\text{per}}^2 \rightarrow \dot{L}_{\text{per}}^2$ is a bounded operator, the bounded perturbation theorem (see [13], Theorem 1.3, on p. 158) implies that the operator

$$A := A_0 + \partial_z^{-1}$$

with the same domain $D(A) = D(A_0) = X_{\text{per}}^1$ also generates a strongly continuous semigroup $(S(t))_{t \geq 0}$ on \dot{L}_{per}^2 . Therefore, the evolution equation in the Cauchy problem (3.21) can be viewed as a bounded perturbation of the evolution equation in the Cauchy problem (3.11). The assertion of the Lemma then follows by Proposition 6.2 on p. 145 in [13]. \square

In what follows, we obtain bounds on the global solution $v \in C(\mathbb{R}, X_{\text{per}}^1)$ to the Cauchy problem (3.21). First, we note the following upper bound on the growth of the global solution.

LEMMA 5. *A global solution $v \in C(\mathbb{R}, X_{\text{per}}^1)$ to the Cauchy problem (3.21) in Lemma 4 satisfies the upper bound*

$$(3.22) \quad \|v(t)\|_{L_{\text{per}}^2} \leq \|v_0\|_{L_{\text{per}}^2} e^{\pi t/6}, \quad t > 0.$$

Proof. Note that the following integration yields

$$\int_{-\pi}^{\pi} v(\partial_z^{-1}v)dz = \frac{1}{2}(\partial_z^{-1}v)^2|_{z=-\pi}^{z=\pi} = 0,$$

since $\partial_z^{-1}v \in H_{\text{per}}^1$ and hence $\partial_z^{-1}v \in C_{\text{per}}$ by Sobolev’s embedding. Integrating by parts yields the following balance equation:

$$\frac{d}{dt} \frac{1}{2} \|v(t)\|_{L_{\text{per}}^2}^2 = \frac{1}{6} \int_{-\pi}^{\pi} v\partial_z [(\pi^2 - z^2)v] dz = -\frac{1}{6} \int_{-\pi}^{\pi} (\pi^2 - z^2)v\partial_z v dz = -\frac{1}{6} \int_{-\pi}^{\pi} z v^2 dz.$$

Hence

$$\frac{d}{dt} \|v(t)\|_{L^2_{\text{per}}}^2 \leq \frac{\pi}{3} \|v(t)\|_{L^2_{\text{per}}}^2$$

and Gronwall's inequality yields the desired bound (3.22). \square

In order to obtain the lower bound on the L^2 norm of the global solution to the Cauchy problem (3.21), we use the generalized method of characteristics and treat $\partial_z^{-1}v(z, t)$ as a source term in (3.11). This term satisfies the following useful bound (also proved in [32]).

LEMMA 6. *If $g := \partial_z^{-1}v \in \dot{H}^1_{\text{per}}$, then*

$$(3.23) \quad \|g\|_{L^\infty_{\text{per}}} \leq \|v\|_{L^1_{\text{per}}}.$$

Proof. By Sobolev embedding of H^1_{per} into C_{per} , g is a continuous 2π -periodic function with zero mean. Therefore, there exists $\zeta \in [-\pi, \pi]$ such that $g(\zeta) = 0$. For every $z \in [-\pi, \pi]$, we can write

$$g(z) = \int_{\zeta}^z v(z') dz',$$

from which bound (3.23) follows. Note that L^2 is continuously embedded into L^1 because of the bound (3.19). \square

By using the family of characteristic curves $z = Z(s, t)$ with $s \in [-\pi, \pi]$ and $t \geq 0$, where Z is defined by the same initial-value problem (3.12), and setting $V(s, t) := v(Z(s, t), t)$ and $G(s, t) := g(Z(s, t), t)$, we obtain the evolution problem in the form

$$(3.24) \quad \begin{cases} \frac{d}{dt} V(s, t) = -\frac{1}{3} Z(s, t) V(s, t) + G(s, t), & t > 0, \\ V(s, 0) = v_0(s). \end{cases}$$

The family of characteristic curves Z is still given by the same explicit form (3.14). Integrating the differential equation (3.24) with an integrating factor yields the explicit solution for V in the form

$$(3.25) \quad V(s, t) = \left[v_0(s) + \int_0^t G(s, t') e^{\frac{1}{3} \int_0^{t'} Z(s, t'') dt''} dt' \right] e^{-\frac{1}{3} \int_0^t Z(s, t') dt'}.$$

By using the explicit solution (3.25), we are able to prove the linear instability result for the Cauchy problem (3.21).

LEMMA 7. *There exists $v_0 \in X^1_{\text{per}}$ and $C > 0$ such that the unique global solution $v \in C(\mathbb{R}, X^1_{\text{per}})$ to the Cauchy problem (3.21) in Lemma 4 satisfies the lower bound*

$$(3.26) \quad \|v(t)\|_{L^2_{\text{per}}} \geq C \|v_0\|_{L^2_{\text{per}}} e^{\pi t/6}, \quad t > 0.$$

Proof. By the chain rule, the explicit expression (3.25) with the help of (3.15) yields the following equation:

$$\begin{aligned} \int_{-\pi}^{\pi} v(z, t)^2 dz &= \int_{-\pi}^{\pi} V(s, t)^2 \frac{\partial Z}{\partial s} ds \\ &= \frac{1}{\pi^2} \int_{-\pi}^{\pi} [\pi \cosh(\pi t/6) - s \sinh(\pi t/6)]^2 \\ &\quad \times \left[v_0(s) + \int_0^t \frac{\pi^2 G(s, t')}{[\pi \cosh(\pi t'/6) - s \sinh(\pi t'/6)]^2} dt' \right]^2 ds. \end{aligned}$$

Let us assume the same constraint $\int_{-\pi}^{\pi} s v_0(s)^2 ds = 0$ as in Lemma 3. Neglecting positive terms in the lower bound, we obtain

$$(3.27) \quad \|v(t)\|_{L^2_{\text{per}}}^2 \geq \frac{1}{4} e^{\pi t/3} \|v_0\|_{L^2_{\text{per}}}^2 - 2 \int_{-\pi}^{\pi} \int_0^t |v_0(s)| |G(s, t')| \frac{[\pi \cosh(\pi t/6) - s \sinh(\pi t/6)]^2}{[\pi \cosh(\pi t'/6) - s \sinh(\pi t'/6)]^2} dt' ds.$$

Let us define for any $t > 0$,

$$K(t, t', s) := \frac{\pi \cosh(\pi t/6) - s \sinh(\pi t/6)}{\pi \cosh(\pi t'/6) - s \sinh(\pi t'/6)}, \quad t' \in [0, t], \quad s \in [-\pi, \pi].$$

We prove that for every $0 \leq t' \leq t$,

$$(3.28) \quad \sup_{s \in [-\pi, \pi]} K(t, t', s) = e^{\pi(t-t')/6}.$$

Indeed, $K(t, t', s) = e^{\pi(t-t')/6} M(t, t', s)$, where

$$M(t, t', s) := \frac{(\pi - s) + (\pi + s)e^{-\pi t/3}}{(\pi - s) + (\pi + s)e^{-\pi t'/3}},$$

and M is monotonically decreasing since $\partial_s M(t, t', s) \leq 0$ for every $t' \in [0, t]$ and $s \in [-\pi, \pi]$. Therefore, M has a maximum at $s = -\pi$, where $M(t, t', -\pi) = 1$.

By using (3.27) and (3.28), we obtain

$$\begin{aligned} \|v(t)\|_{L^2_{\text{per}}}^2 &\geq \frac{1}{4} e^{\pi t/3} \|v_0\|_{L^2_{\text{per}}}^2 - 2 \|v_0\|_{L^1_{\text{per}}} \int_0^t \|g(t')\|_{L^\infty_{\text{per}}} e^{\pi(t-t')/3} dt' \\ &\geq \frac{1}{4} e^{\pi t/3} \|v_0\|_{L^2_{\text{per}}}^2 - 2 \|v_0\|_{L^1_{\text{per}}} \int_0^t \|v(t')\|_{L^1_{\text{per}}} e^{\pi(t-t')/3} dt' \\ &\geq \frac{1}{4} e^{\pi t/3} \|v_0\|_{L^2_{\text{per}}}^2 - 2\sqrt{2\pi} \|v_0\|_{L^1_{\text{per}}} \|v_0\|_{L^2_{\text{per}}} e^{\pi t/3} \int_0^t e^{-\pi t'/6} dt', \end{aligned}$$

where (3.20), (3.22), and (3.23) have been used in the last two inequalities. Hence,

$$\|v(t)\|_{L^2_{\text{per}}}^2 e^{-\pi t/3} \geq \|v_0\|_{L^2_{\text{per}}}^2 \left(\frac{1}{4} \|v_0\|_{L^2_{\text{per}}}^2 - \frac{12\sqrt{2}}{\sqrt{\pi}} \|v_0\|_{L^1_{\text{per}}} \right)$$

and since $\|v_0\|_{L^2_{\text{per}}}$ can be much larger than $\|v_0\|_{L^1_{\text{per}}}$ by the bound (3.19), there exist $v_0 \in X^1_{\text{per}}$ and $C^2 \in (0, 1/4)$ such that

$$(3.29) \quad \|v_0\|_{L^1_{\text{per}}} \leq \frac{\sqrt{\pi}(1 - 4C^2)}{48\sqrt{2}} \|v_0\|_{L^2_{\text{per}}},$$

and hence

$$(3.30) \quad \|v(t)\|_{L^2_{\text{per}}}^2 e^{-\pi t/3} \geq C^2 \|v_0\|_{L^2_{\text{per}}}^2.$$

This yields the desired bound (3.26). □

Remark 15. Let us show that there exist functions $v_0 \in X_{\text{per}}^1$ satisfying the constraints (3.8), (3.9), and (3.29). Indeed, if v_0 is odd, then v_0^2 is even, and hence the two constraints (3.8) and (3.9) are satisfied simultaneously. From the class of odd initial data we need to pick functions in X_{per}^1 that satisfy the inequality (3.29) for a fixed $C^2 \in (0, 1/4)$. For example, we can consider the following odd function in $\dot{H}_{\text{per}}^1 \subset X_{\text{per}}^1$:

$$(3.31) \quad v_0(x) = \frac{x(\pi^2 - x^2)}{1 + a^2x^2}, \quad x \in [-\pi, \pi],$$

where $a > 0$ is a parameter. We obtain by direct computation

$$\|v_0\|_{L_{\text{per}}^1} = \left(\frac{\pi^2}{a^2} + \frac{1}{a^4}\right) \log(1 + \pi^2 a^2) - \frac{\pi^2}{a^2}$$

and

$$\|v_0\|_{L_{\text{per}}^2}^2 = \frac{1}{a^3} \left[\left(\pi^4 + \frac{6\pi^2}{a^2} + \frac{5}{a^4}\right) \arctan(\pi a) - \frac{\pi(15 + 13\pi^2 a^2)}{3a^3} \right].$$

Since $\|v_0\|_{L_{\text{per}}^1} = \mathcal{O}(\log(a)a^{-2})$ decays to zero as $a \rightarrow \infty$ faster than $\|v_0\|_{L_{\text{per}}^2} = \mathcal{O}(a^{-3/2})$, inequality (3.29) can be satisfied for sufficiently large a .

Remark 16. If $v_0(\pm\pi) = 0$ like in the example (3.31), then $v_0 \in \dot{H}_{\text{per}}^1$ and the truncated linearized evolution (3.11) preserves the constraint $v(\pm\pi, t) = 0$ for every $t \in \mathbb{R}$ (see Remark 9). However, the integral term $\partial_z^{-1}v$ in the full linearized evolution (3.21) does not generally preserve the same constraint because it is uniquely defined from the condition that $\partial_z^{-1}v$ has zero mean. As a result, the full linearized equation does not generally admit a solution $v \in C(\mathbb{R}, \dot{H}_{\text{per}}^1)$ even if $v_0 \in \dot{H}_{\text{per}}^1$.

Remark 17. In the presence of the source term G , we are not able to show that $\|v(t)\|_{L_{\text{per}}^1}$ remains bounded as $t \rightarrow \infty$ (see Remark 11). By using the integral

$$\int_{-\pi}^{\pi} \frac{\pi^2}{[\pi \cosh(\pi t'/6) - s \sinh(\pi t'/6)]^2} ds = 2\pi, \quad t' \in [0, t],$$

we obtain the bound

$$\|v(t)\|_{L_{\text{per}}^1} \leq \|v_0\|_{L_{\text{per}}^1} + 2\pi \int_0^t \|g(t')\|_{L_{\text{per}}^\infty} dt',$$

in view of (3.15) and (3.25). Thanks to the bound (3.23), the inequality is closed as follows:

$$\|v(t)\|_{L_{\text{per}}^1} \leq \|v_0\|_{L_{\text{per}}^1} + 2\pi \int_0^t \|v(t')\|_{L_{\text{per}}^1} dt'.$$

By Gronwall’s inequality, this bound gives the fast exponential growth

$$\|v(t)\|_{L_{\text{per}}^1} \leq \|v_0\|_{L_{\text{per}}^1} e^{2\pi t},$$

which cannot be sharp because $\|v(t)\|_{L_{\text{per}}^1}$ is bounded by a slowly growing exponential function that follows from the bounds (3.20) and (3.22).

Remark 18. There exists a conserved energy for the Cauchy problem (3.21) (see Remark 12) which is given by

$$(3.32) \quad \langle Lv(t), v(t) \rangle = \langle Lv_0, v_0 \rangle,$$

where the self-adjoint operator L is defined by (3.2). However, the conserved quantity (3.32) does not prevent $\|v(t)\|_{L_{\text{per}}^2}$ from growing exponentially fast as $t \rightarrow \infty$ because the bounded operator L is not coercive under the constraint (3.9) (see Lemma 8).

3.3. Spectrum of the linear self-adjoint operator L . Here we consider the spectrum $\sigma(L)$ of the linear self-adjoint operator L defined by (3.2). We will prove that $\sigma(L)$ consists of the continuous spectrum on $[0, \pi^2/6]$, which includes the embedded eigenvalue $\lambda_0 = 0$ with the eigenvector $\partial_z U$ and a simple negative eigenvalue $\lambda_1 < 0$. No spectral gap appears between $\lambda_0 = 0$ and the continuous spectrum. The following lemma gives the corresponding result.

LEMMA 8. *The spectrum of the self-adjoint operator L given by (3.2) is*

$$(3.33) \quad \sigma(L) = \{\lambda_1\} \cup \left[0, \frac{\pi^2}{6}\right],$$

where $\lambda_1 < 0$ is the unique zero of the transcendental equation

$$(3.34) \quad (\pi^2 + 3\lambda) \log \frac{\sqrt{\pi^2 - 6\lambda} + \pi}{\sqrt{\pi^2 - 6\lambda} - \pi} - 3\pi\sqrt{\pi^2 - 6\lambda} = 0, \quad \lambda < 0.$$

Proof. By the spectral theorem (see, e.g., Definition 8.39 and Theorems 8.70 and 8.71 in [35]), the spectrum of the self-adjoint operator L in \dot{L}^2_{per} denoted by $\sigma(L)$ may consist of only two disjoint sets on the real line: the point spectrum of eigenvalues with eigenvectors in \dot{L}^2_{per} denoted by $\sigma_p(L)$ and the continuous spectrum denoted by $\sigma_c(L)$, where the resolvent operator exists but is unbounded.

The self-adjoint operator L in (3.2) is given by the sum of a bounded operator L_0 and a compact operator K given by

$$(3.35) \quad L_0 := \frac{1}{6}P_0(\pi^2 - z^2)P_0 : \dot{L}^2_{\text{per}} \rightarrow \dot{L}^2_{\text{per}}$$

and

$$(3.36) \quad K := P_0\partial_z^{-2}P_0 : \dot{L}^2_{\text{per}} \rightarrow \dot{L}^2_{\text{per}}.$$

Moreover, the compact operator is in the trace class since $\sum_{n=1}^\infty n^{-2} < \infty$. By Kato's theorem [27] (see Theorem 4.4 on p. 540 in [28]), $\sigma_c(L) = \sigma_c(L_0)$. We show that $[0, \pi^2/6] \subseteq \sigma_c(L_0)$ by considering the odd functions in \dot{L}^2_{per} , which can be represented by the Fourier sine series. Let us denote the space of odd functions in \dot{L}^2_{per} by $L^2_{\text{per,odd}}$. Then,

$$L_0f = \frac{1}{6}(\pi^2 - z^2)f \quad \forall f \in L^2_{\text{per,odd}}.$$

Then, $\sigma_c(L_0)$ in $L^2_{\text{per,odd}}$ coincides with the range of the multiplicative function $h(z) = \frac{1}{6}(\pi^2 - z^2)$ for $z \in [-\pi, \pi]$, which is $[0, \pi^2/6]$. Hence, $[0, \pi^2/6] \subseteq \sigma_c(L_0)$ in \dot{L}^2_{per} .

Let us show that $[0, \pi^2/6] \equiv \sigma_c(L_0)$ by working with the resolvent equation $(L_0 - \lambda I)f = g$ for given $g \in \dot{L}^2_{\text{per}}$ and $\lambda \notin [0, \pi^2/6]$. The resolvent equation can be written in the component form for $z \in [-\pi, \pi]$:

$$\frac{1}{6}(\pi^2 - 6\lambda - z^2)f(z) - k(f) = g(z), \quad k(f) := \frac{1}{12\pi} \int_{-\pi}^\pi (\pi^2 - z^2)f(z)dz,$$

where $f \in \dot{L}^2_{\text{per}}$ is supposed to satisfy the zero-mean constraint $\int_{-\pi}^\pi f(z)dz = 0$. Computing the solution explicitly,

$$f(z) = \frac{6}{\pi^2 - 6\lambda - z^2} [g(z) + k(f)],$$

and using the zero mean constraint, we can define $k(f)$ in terms of g :

$$k(f) = \frac{\int_{-\pi}^{\pi} \frac{g(z)}{\pi^2 - 6\lambda - z^2} dz}{\int_{-\pi}^{\pi} \frac{1}{\pi^2 - 6\lambda - z^2} dz}.$$

For every $\lambda \notin [0, \pi^2/6]$, there exist positive constants $C_\lambda, C'_\lambda > 0$ such that

$$\sup_{z \in [-\pi, \pi]} \frac{6}{|\pi^2 - 6\lambda - z^2|} \leq C_\lambda, \quad \left| \int_{-\pi}^{\pi} \frac{6}{\pi^2 - 6\lambda - z^2} dz \right| \geq C'_\lambda.$$

As a result, we obtain the bound

$$\|f\|_{L^2_{\text{per}}} \leq C_\lambda \left[\|g\|_{L^2_{\text{per}}} + |k(f)|\sqrt{2\pi} \right] \leq C_\lambda [1 + 2\pi(C'_\lambda)^{-1}C_\lambda] \|g\|_{L^2_{\text{per}}}.$$

Therefore, the resolvent operator $(L_0 - \lambda I)^{-1} : \dot{L}^2_{\text{per}} \rightarrow \dot{L}^2_{\text{per}}$ is bounded for every $\lambda \notin [0, \pi^2/6]$ so that $\sigma_c(L_0) = [0, \pi^2/6]$.

In order to study $\sigma_p(L) \in \mathbb{R} \setminus [0, \pi^2/6]$, we consider the spectral problem for operator L with the spectral parameter $\lambda \notin [0, \pi^2/6]$:

$$(3.37) \quad \frac{1}{6}P_0(\pi^2 - z^2)w + P_0\partial_z^{-2}w = \lambda w, \quad w \in \dot{L}^2_{\text{per}}.$$

Since $\partial_z^{-2}w \in H^2_{\text{per}}$, bootstrapping arguments show that $w \in H^2_{\text{loc}}$ on any compact subset in $(-\pi, \pi)$. Iterations of bootstrapping arguments yield $w \in H^\infty_{\text{loc}}$. Therefore, the spectral problem (3.37) can be differentiated twice on a compact subset in $(-\pi, \pi)$, after which it is rewritten as the second-order differential equation

$$(3.38) \quad (\pi^2 - z^2 - 6\lambda) \frac{d^2w}{dz^2} - 4z \frac{dw}{dz} + 4w(z) = 0, \quad w \in H^\infty_{\text{loc}},$$

with the two linearly independent solutions for $\lambda \in \mathbb{R} \setminus [0, \pi^2/6]$,

$$w_1(z) = z$$

and

$$w_2(z) = \begin{cases} -1 + \frac{z^2}{2(\pi^2 - z^2 - 6\lambda)} + \frac{3z}{4\sqrt{\pi^2 - 6\lambda}} \log \frac{\sqrt{\pi^2 - 6\lambda} + z}{\sqrt{\pi^2 - 6\lambda} - z}, & \lambda < 0, \\ -1 + \frac{z^2}{2(\pi^2 - z^2 - 6\lambda)} - \frac{3z}{2\sqrt{6\lambda - \pi^2}} \arctan \frac{z}{\sqrt{6\lambda - \pi^2}}, & \lambda > \frac{\pi^2}{6}. \end{cases}$$

The first solution corresponds to the eigenvector $\partial_z U$ of the spectral problem (3.37) for the eigenvalue $\lambda_0 = 0$, which is embedded into $\sigma_c(L) = [0, \pi^2/6]$. Since eigenvectors of the self-adjoint operator for distinct eigenvalues are orthogonal, we are looking for solutions w of the spectral problem (3.37) such that $\langle w, w_1 \rangle = 0$. Therefore, we take⁴ $w = w_2$ and extend it from H^2_{loc} to \dot{L}^2_{per} . This extension is achieved if and only if w has zero mean, that is,

$$(3.39) \quad 0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} w_2(z) dz = \begin{cases} -\frac{3}{4} + \frac{\pi^2 + 3\lambda}{4\pi\sqrt{\pi^2 - 6\lambda}} \log \frac{\sqrt{\pi^2 - 6\lambda} + \pi}{\sqrt{\pi^2 - 6\lambda} - \pi}, & \lambda < 0, \\ -\frac{3}{4} - \frac{\pi^2 + 3\lambda}{2\pi\sqrt{6\lambda - \pi^2}} \arctan \frac{\pi}{\sqrt{6\lambda - \pi^2}}, & \lambda > \frac{\pi^2}{6}. \end{cases}$$

The piecewise graph of the right-hand side of the zero-mean constraint (3.39) on $(-\infty, 0)$ and $(\pi^2/6, \infty)$ is shown in Figure 3. The first line of the zero-mean constraint (3.39) is equivalent to the transcendental equation (3.34) and it has only one simple zero at $\lambda_1 \approx -0.2262$. The second line of (3.39) does not have any zeros. Hence, $\lambda_1 < 0$ is the only eigenvalue in $\sigma_p(L)$. \square

⁴Note that $\langle w_2, w_1 \rangle = 0$ because w_1 is odd and w_2 is even.




FIG. 3. The graph of the right-hand side of the zero-mean constraint (3.39) on $(-\infty, 0)$ and $(\pi^2/6, \infty)$ as a function of the spectral parameter λ . Only one simple zero $\lambda_1 < 0$ exists.

Remark 19. For the smooth periodic waves of Lemma 1, we proved in [18] that $\sigma(L)$ in \dot{L}_{per}^2 includes a simple negative eigenvalue and a simple zero eigenvalue with the eigenvector $\partial_z U$ and the rest of the spectrum is positive and bounded away from zero. Hence, the spectral gap is present in the case of smooth periodic waves. This enabled us in [18] to use the Hamilton–Krein index theory to deduce that $\sigma(\partial_z L) \subset i\mathbb{R}$ and hence to deduce the spectral stability of the smooth periodic waves according to Definition 2. By the standard analysis involving the conserved quantity (3.32) (see [23]), this spectral stability result transfers to the linear stability of the smooth periodic waves according to Definition 3. In the spectral problem for the peaked periodic wave, however, this spectral gap is not present. Therefore, we are not able to deduce spectral instability of the peaked periodic wave from the spectrum of L .

4. Discussion. We have studied peaked periodic traveling wave solutions of the reduced Ostrovsky equation (1.1). We found that the peaked periodic wave with parabolic shape U_* is the unique peaked periodic traveling wave with a single minimum per period and that the boundary-value problem (2.1) does not admit Hölder continuous solutions (see Lemma 2). As a consequence, the existence of cusped waves obtained in [40] as well as the existence of small-amplitude peaked waves found in [21] was disproved.

Furthermore, we proved that the peaked periodic wave U_* is linearly unstable with respect to co-periodic perturbations in the space X_{per}^1 , which is the maximal domain of the linearized operator $\partial_z L$ (see Lemma 7). This result was obtained using sharp exponential bounds on the L^2 norm of perturbations v of U_* in X_{per}^1 satisfying the Cauchy problem (3.1) with the peaked periodic wave U_* and for the wave speed c_* .

Passing from linear to nonlinear instability is often a delicate issue. Several authors have shown that linear instability directly implies nonlinear instability if a part of the spectrum of the linearized operator is located in the right half of the complex plane: see, for instance, [17, 37] and Theorem 5.1.5 in [25]. However, these approaches do not work for the reduced Ostrovsky equation since the linearized evolution is defined in the space X_{per}^1 whereas the nonlinear evolution is defined in \dot{H}_{per}^s with $s > 3/2$.

It is not clear if the local well-posedness results can be extended to the space X_{per}^1 . It is also unclear how the peaks of the peaked periodic wave move under the flow of the reduced Ostrovsky equation (1.1). For these reasons, the nonlinear instability of the peaked periodic wave in the reduced Ostrovsky equation remains an open problem for now.

Acknowledgments. This project was initiated during the research program on Nonlinear Water Waves at the Isaac Newton Institute at Cambridge in August 2017. Both authors thank Mats Ehrnström for useful discussions during the program and afterward. Computations in the proof of Lemma 8 were performed back in 2015 in collaboration with Ted Johnson (UCL). Finally, the authors thank the referees for useful remarks which helped to improve the manuscript.

REFERENCES

- [1] A. BOUTET DE MONVEL AND D. SHEPELSKY, *The Ostrovsky–Vakhnenko equation by a Riemann–Hilbert approach*, J. Phys. A, 48 (2015) 035204.
- [2] A. BRESSAN, *Hyperbolic Systems of Conservation Laws: The One-Dimensional Cauchy Problem*, Oxford Lecture Ser. Math. Appl. 20, Oxford University Press, Oxford, 2000.
- [3] J. C. BRONSKI, M. A. JOHNSON, AND T. KAPUTULA, *An index theorem for the stability of periodic traveling waves of KdV Type*, Proc. Roy. Soc. Edinburgh Sect. A, 141 (2011), pp. 1141–1173.
- [4] G. BRUELL AND R. N. DHARA, *Waves of Maximal Height for a Class of Nonlocal Equations with Homogeneous Symbol*, preprint, <https://arxiv.org/abs/1810.00248v1>, 2018.
- [5] J. C. BRUNELLI AND S. SAKOVICH, *Hamiltonian structures for the Ostrovsky–Vakhnenko equation*, Commun. Nonlinear Sci. Numer. Simul., 18 (2013), pp. 56–62.
- [6] G. M. COCLITE AND L. DIRUVO, *Convergence of the Ostrovsky equation to the Ostrovsky–Hunter one*, J. Differential Equations, 256 (2014), pp. 3245–3277.
- [7] G. M. COCLITE AND L. DIRUVO, *Oleinik type estimates for the Ostrovsky–Hunter equation*, J. Math. Anal. Appl., 423 (2015), pp. 162–190.
- [8] G. M. COCLITE AND L. DIRUVO, *Well-posedness of bounded solutions of the non-homogeneous initial-boundary value problem for the Ostrovsky–Hunter equation*, J. Hyperbolic Differ. Equ., 12 (2015), pp. 221–248.
- [9] A. CONSTANTIN AND W. A. STRAUSS, *Stability of peakons*, Comm. Pure Appl. Math., 53 (2000), pp. 603–610.
- [10] M. EHRNSTRÖM, M. JOHNSON, AND K. M. CLAASSEN, *Existence of a highest wave in a fully dispersive two-wave shallow water model*, Arch. Ration. Mech. Anal. (2018), <https://doi.org/10.1007/s00205-018-1306-5>.
- [11] M. EHRNSTRÖM AND H. KALISCH, *Global Bifurcation for the Whitham Equation*, Math. Model. Nat. Phenom., 8 (2013), pp. 13–30.
- [12] M. EHRNSTRÖM AND E. WAHLÉN, *On Whitham’s conjecture of a highest cusped wave for a nonlocal shallow water wave equation*, Ann. Inst. H. Poincaré Anal. Non Linéaire, to appear.
- [13] K. J. ENGEL AND R. NAGEL, *One-Parameter Semigroups for Linear Evolution Equations*, Springer-Verlag, Berlin, 2000.
- [14] E. R. JOHNSON AND D. E. PELINOVSKY, *Orbital stability of periodic waves in the class of reduced Ostrovsky equations*, J. Differential Equations, 261 (2016), pp. 3268–3304.
- [15] B. F. FENG, K. MARUNO, AND Y. OHTA, *On the τ -functions of the reduced Ostrovsky equation and the $A_2^{(2)}$ two-dimensional Toda system*, J. Phys. A, 45 (2012), 355203.
- [16] B. F. FENG, K. MARUNO, AND Y. OHTA, *Integrable semi-discretizations of the reduced Ostrovsky equation*, J. Phys. A, 48 (2015), 135203.
- [17] S. FRIEDLANDER, W. STRAUSS, AND M. VISHIK, *Nonlinear instability in an ideal fluid*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 14 (1997), pp. 187–209.
- [18] A. GEYER AND D. E. PELINOVSKY, *Spectral stability of periodic waves in the generalized reduced Ostrovsky equation*, Lett. Math. Phys., 107 (2017), pp. 1293–1314.
- [19] R. H. J. GRIMSHAW, K. HELFRICH, AND E. R. JOHNSON, *The reduced Ostrovsky equation: Integrability and breaking*, Stud. Appl. Math., 129 (2012), pp. 414–436.
- [20] R. GRIMSHAW AND D. E. PELINOVSKY, *Global existence of small-norm solutions in the reduced Ostrovsky equation*, Discrete Contin. Dyn. Syst. A, 34 (2014), pp. 557–566.

- [21] S. HAKKAEV, M. STANISLAVOVA, AND A. STEFANOV, *Periodic travelling waves of the regularized short pulse and Ostrovsky equations: Existence and stability*, SIAM J. Math. Anal., 49 (2017), pp. 674–698.
- [22] S. HAKKAEV, M. STANISLAVOVA, AND A. STEFANOV, *Spectral stability for classical periodic waves of the Ostrovsky and short pulse models*, Stud. Appl. Math., 139 (2017), pp. 405–433.
- [23] M. HARAGUS, J. LI, AND D. E. PELINOVSKY, *Counting unstable eigenvalues in Hamiltonian spectral problems via commuting operators*, Comm. Math. Phys., 354 (2017), pp. 247–268.
- [24] M. HARAGUS AND T. KAPITULA, *On the spectra of periodic waves for infinite-dimensional Hamiltonian systems*, Phys. D, 237 (2008), pp. 2649–2671.
- [25] D. HENRY, *Geometric Theory of Semilinear Parabolic Equations*, Springer-Verlag, Berlin, 1981.
- [26] J. K. HUNTER, *Numerical solution of some nonlinear dispersive wave equations*, in Computational Solution of Nonlinear Systems of Equations, E. L. Allgower and K. Georg, eds., Lectures in Appl. Math. 26, AMS, Providence, RI, 1990, pp. 301–316.
- [27] T. KATO, *Perturbation of continuous spectra by trace class operators*, Proc. Japan Acad., 33 (1957), pp. 260–264.
- [28] T. KATO, *Perturbation Theory for Linear Operators*, Springer-Verlag, Berlin, 1995.
- [29] R. A. KRAENKEL, H. LEBLOND, AND M. A. MANNA, *An integrable evolution equation for surface waves in deep water*, J. Phys. A, 47 (2014), 025208.
- [30] J. LENELLS, *Stability of periodic peakons*, Int. Math. Res. Not. IMRN, 2004, pp. 485–499.
- [31] J. LENELLS, *Stability for the periodic Camassa-Holm equation*, Math. Scand., 97 (2005), pp. 188–200.
- [32] Y. LIU, D. PELINOVSKY, AND A. SAKOVICH, *Wave breaking in the Ostrovsky–Hunter equation*, SIAM J. Math. Anal., 42 (2010), pp. 1967–1985.
- [33] L. A. OSTROVSKY, *Nonlinear internal waves in a rotating ocean*, Okeanologia, 18 (1978), pp. 181–191.
- [34] D. E. PELINOVSKY, *Spectral stability of nonlinear waves in KdV-type evolution equations*, in Nonlinear Physical Systems: Spectral Analysis, Stability, and Bifurcations, O. N. Kirillov and D. E. Pelinovsky, eds., Wiley, New York, 2014, pp. 377–400.
- [35] M. RENARDY AND R. C. ROGERS, *An Introduction to Partial Differential Equations*, 2nd ed., Texts Appl. Math. 13, Springer-Verlag, New York, 2004.
- [36] N. SANFORD, K. KODAMA, J. D. CARTER, AND H. KALISCH, *Stability of traveling wave solutions to the Whitham equation*, Phys. Lett. A., 378 (2014), pp. 2100–2107.
- [37] J. SHATAH AND W. STRAUSS, *Spectral condition for instability*, Contemp. Math., 255 (2000), pp. 189–198.
- [38] M. STANISLAVOVA AND A. STEFANOV, *On the spectral problem $Lu = \lambda u'$ and applications*, Comm. Math. Phys., 343 (2016), pp. 361–391.
- [39] A. STEFANOV, Y. SHEN, AND P. G. KEVREKIDIS, *Well-posedness and small data scattering for the generalized Ostrovsky equation*, J. Differential Equations, 249 (2010), pp. 2600–2617.
- [40] YU. A. STEPANYANTS, *On stationary solutions of the reduced Ostrovsky equation: Periodic waves, compactons and compound solitons*, Chaos, 28 (2006), pp. 193–204.
- [41] G. STOKES, *On the theory of oscillatory waves*, J. Math. Phys. Pap., 1 (1847), pp. 314–326.
- [42] J. TOLAND, *On the existence of a wave of greatest height and Stokes’s conjecture*, Proc. Roy. Soc. London, Ser. A, 363 (1978), pp. 469–485.
- [43] V. A. VAKHNENKO, *Solitons in a nonlinear model medium*, J. Phys. A, 25 (1992), pp. 4181–4187.
- [44] V. A. VAKHNENKO AND E. J. PARKES, *The calculation of multi-soliton solutions of the Vakhnenko equation by the inverse scattering method*, Chaos Solitons Fractals, 13 (2002), pp. 1819–1826.